REMARKS

Entry of the foregoing and favorable reconsideration and reexamination of the subject matter pursuant to and consistent with 37 C.F.R. § 1.112 is respectfully requested.

Applicant requests consideration of these additional remarks which supplement the response file on June 29, 1999. It will be demonstrated below that the current claims of record are clearly novel and unobvious over the cited prior art of record and should be taken into consideration by the Examiner.

It should be recalled that the present claims of record are directed to a method aiming at inhibiting the replication of an immunodeficiency retrovirus wherein 100% inhibition of the retrovirus in primary cultures of monocytes in the host is achieved using selected muramyl peptides.

It should be brought to the immediate attention of the Examiner that:

- (1) the claims of record encompass 100% inhibition of the retrovirus which is not disclosed in the prior art; and
- (2) this inhibition was demonstrated in primary cultures (cultures prepared directly from the tissues of an organism) of monocytes of the host, which is also not demonstrated in the cited prior art.

Applicant submits that the demonstration in the present invention of 100% inhibition of a retrovirus in primary cultures of monocytes is an extremely important aspect of the present invention that must be taken into

consideration by the Examiner in analyzing the prior art. This is because it is known in the art that the use of primary cultures of monocytes is a more scientifically sound *in vitro* system for testing drugs or medicaments for the inhibition of HIV-1 than in those cell lines disclosed in the prior art, as will be discussed more extensively below under the heading HIV-1 replication.

It should be emphasized, as will be discussed in greater detail below, that the cited prior art teaches the use of muramyl peptides for inhibiting HIV-1 infection using strains that are infected by T-Tropic HIV-1 strains. The prior art is silent with respect to the use of muramyl peptides for inhibiting immunodeficiency retroviruses in the presently claimed primary cultures of monocytes which are infected by M-Tropic HIV-1 strains.

#### HIV-1 REPLICATION

It is now known that HIV-1 needs to replicate in macrophages or dendritic cells prior to spreading to T lymphocytes. At the early stages of HIV-1 infection, shortly after seroconversion and during the asymptomatic period of AIDS, macrophage tropic or M-Tropic strains of the virus predominate.

In contrast, in the late stages of HIV-1 disease in association with CD4 T cell decline and progression to AIDS, T cell lines or T-Tropic strains of HIV-1 predominate.

The mechanism behind entry of HIV-1 gp120 at the different stages of HIV-1 disease is different. It is now known that besides binding to the CD4 receptor, interaction of the V3 loop in gp120 with a second receptor or co-receptor is required for gp120 to enter the cells. At the early stages of HIV-1 disease the co-receptor required for the gp120 to enter the macrophages

consideration by the Examiner in analyzing the prior art. This is because it is known in the art that the use of primary cultures of monocytes is a more scientifically sound *in vitro* system for testing drugs or medicaments for the inhibition of HIV-1 than in those cell lines disclosed in the prior art, as will be discussed more extensively below under the heading HIV-1 replication.

It should be emphasized, as will be discussed in greater detail below, that the cited prior art teaches the use of muramyl peptides for inhibiting HIV-1 infection using strains that are infected by T-Tropic HIV-1 strains. The prior art is silent with respect to the use of muramyl peptides for inhibiting immunodeficiency retroviruses in the presently claimed primary cultures of monocytes which are infected by M-Tropic HIV-1 strains.


#### HIV-1 REPLICATION

It is now known that HIV-1 needs to replicate in macrophages or dendritic cells prior to spreading to T lymphocytes. At the early stages of HIV-1 infection, shortly after seroconversion and during the asymptomatic period of AIDS, macrophage tropic or M-Tropic strains of the virus predominate.

In contrast, in the late stages of HIV-1 disease in association with CD4 T cell decline and progression to AIDS, T cell lines or T-Tropic strains of HIV-1 predominate.

The mechanism behind entry of HIV-1 gp120 at the different stages of HIV-1 disease is different. It is now known that besides binding to the CD4 receptor, interaction of the V3 loop in gp120 with a second receptor or co-receptor is required for gp120 to enter the cells. At the early stages of HIV-1 disease the co-receptor required for the gp120 to enter the macrophages

was discovered to be the CCR-5 co-receptor. In contrast, in the late stages of the disease, the co-receptor required to enter the cells was discovered to be fusin or the CXCR-4 co-receptor. These receptors are different as can be seen schematically below:



Both of the co-receptors were discovered to be chemokine receptors that belong to the family of G-coupled protein receptors, which have seven transmembrane regions. The fact that these receptors have seven transmembrane regions is important, since resistance to HIV-1 infection, including T cell depletion, was discovered in certain individuals bearing a mutant allele of the CCR-5 chemokine co-receptor.

This mutant CCR-5 co-receptor lacks the three transmembrane segments of the wild CCR-5 receptor and was unable to support membrane fusion by both the primary and dual-tropic virus env. Hence, it was concluded that homozygous individuals having this mutant CCR-5 receptor are highly resistant to HIV-1 infection.

The above supports the theory that the CCR-5 receptor plays a primary role in the replication of HIV-1 which replication then leads in the later stages of the disease to AIDS

Hence, it is more scientifically beneficial to find drugs that target the early stages of HIV-1 infection using the M-Tropic HIV-1 strain at an early stage of infection, thus either diminishing or preventing the total onset of AIDS. This even the more so since macrophages serve as a reservoir for the virus and this reservoir is less sensitive to antiretroviral effects than T-lymphocytes.

Therefore, in the present invention inhibition of the replication of HIV-1, in the presently claimed process was demonstrated **in primary cultures of monocytes** (monocytes are precursors to macrophages) which cultures are the scientific tools of choice to use in drug evaluation experiments for HIV-1 inhibition, as explained above.

In contrast the **use of cell lines** to test for drugs which inhibit HIV-1 is highly artificial and drugs that can inhibit T-Tropic HIV replication are not necessarily effective against replication of M-Tropic viruses in macrophages. This has been demonstrated by the fact that SDF-1 (Stromal cell derived factor), the ligand for CXCR-4, can inhibit virus entry into cell lines, but has absolutely no effect of preventing M-Tropic HIV-1 entry and infection in macrophages or primary T-lymphocytes.

Therefore, the fact that the Applicant has demonstrated 100% inhibition of a retrovirus in primary cultures of monocytes (M-Tropic HIV-1 strains) with the presently claimed muramyl peptides is an unexpected result which should distinguish clearly over the prior art of record where such a demonstration is not achieved. Rather the cited prior art teaches low inhibition of several muramyl peptides in cell lines which are T-Tropic strains of HIV-1.

These and other differences will be addressed in view of the issues brought to bear in the last Official Action.

#### 35 U.S.C. §102(b)

The Examiner deems that Claims 14 to 21, 25, 26, 28 to 30 and 34 lack novelty in view of Schreck et al.

Furthermore, Claims 14 to 21, 25, 26, 28 to 30 and 34 lack novelty over Masihi et al.

#### Schreck et al.

Schreck et al. teach the use of muramyl peptides as **adjuvants** in potential vaccines against AIDS. By definition an adjuvant is an ingredient (as in a prescription or solution) that modifies the action of the principle ingredient. An adjuvant is not the active ingredient in a vaccine, as the skilled artisan well knows.

Furthermore, Schreck et al. disclose that it would be beneficial to select adjuvants that do not induce NF- $\kappa$ B activation and particularly if the

vaccines are to be aimed at treating seropositive individuals, since it was believed that the **activation of NF- $\kappa$ B purportedly enhanced HIV-1 expression**.

In fact, MDP (thr)-GDP was found **to be the only lipophilic, nonpyrogenic adjuvant that demonstrated lack of NF- $\kappa$ B activation**. This teaching is apparent at page 188, 2<sup>nd</sup> column, lines 13 to 15 of Schreck et al.

Although two muramyl peptides, encompassed by the present claims were tested for NF- $\kappa$ B activation, it was discovered that in the human Mono-Mac-6 cell line **NF- $\kappa$ B activation was apparent using murabutide and murametide** as set forth in the sentence bridging column 1 and column 2 at page 190 of Schreck et al. Therefore, murametide and murabutide do not belong to the selected category of an adjuvant that could be foreseen for use with an AIDS vaccine.

Furthermore, it is apparent that there is no experimental evidence that the muramyl peptides utilized in Schreck et al. can inhibit the replication of immunodeficiency retroviruses. Thus, a skilled artian can conclude nothing about whether the muramyl peptides in Schreck et al. have any inhibitory properties.

Therefore, Applicant submits that since Schreck et al. fails to teach the use of the claimed muramyl peptides as an active ingredient in a process to inhibit immunodeficiency retroviruses and since the claimed muramyl peptides do not fall into the category of those being sought in Schreck et al., the presently claimed invention is not anticipated by Schreck et al.

**Masihi et al.**

Masihi et al. disclose that muramyl dipeptide can enhance monocyte/macrophage CSF in serum and promote nonspecific resistance against a variety of microbial pathogens including HIV infection of CD4<sup>+</sup> H9 lymphocytes and U937 monocytic cells. However, this effect cannot be mediated by macrophage-CSF which itself has been shown to increase viral replication (see, Annex I, page 33, last paragraph, left column).

The Examiner refers to page 397 of Masihi et al. where murabutide was taught to be used as an **adjuvant** in human clinical trials. As discussed above, an adjuvant is solely used as a vehicle to modify the action of the active ingredient. Masihi et al. fails to teach that murabutide can be used in a process to treat immunodeficiency retroviruses directly.

Indeed, the cell lines used in the experiments in as the active ingredient in the manufacture of a medicament are H9, KE37/1 and U937 which are only infectable by T-Tropic HIV-1 strains. In contrast the present invention uses primary cultures of monocytes which are only infectable by M-Tropic HIV-1 strain. Thus, Masihi et al. disclose muramyl peptides for targeting the late stages of HIV-1, while the muramyl peptides in the process of the presently claimed invention target the early stage of HIV-1.

Therefore, in view of the above, Applicant submits that the presently claimed invention is not anticipated by Masihi et al.

35 U.S.C. §103(a)

Masihi et al.



Masihi et al. fail to teach the skilled artisan that murabutide can be used in a medicament as the active ingredient for inhibiting the replication of a retrovirus. Rather Masihi et al. teach the use of murabutide only as an adjuvant.

Furthermore, a skilled artisan would not extrapolate the results of a muramyl dipeptide disclosed in Masihi et al. to include all muramyl peptides, since as taught in Masihi et al. at page 189 under Reagents, different muramyl peptides have different properties.

Only if the Examiner deems that a skilled artisan would indeed extrapolate results from MDP to the rest of the muramyl peptides, Applicant would like to point out that Masihi et al. discloses only 67% reduction of the p24 antigen using MDP and only a 38% inhibition on day 14 using infected CD4<sup>+</sup> KE37/L lymphocytes and further teaches that 1000 µg/ml dosages were more effective.

Moreover, Figure 3 clearly demonstrates that less than 50% inhibition of p24 antigen using MDP at 1,000 µg/ml is achieved in U937 monocytic cells. **This percentage inhibition cannot be compared to the 100% inhibition achieved by the claimed muramyl compounds of the present invention,** which Applicant submits is an unexpected result.

Furthermore, Masihi et al. teach using 1000 µg/ml MDP which is an extremely high dosage and the side effects of MDP, including pyrogenicity and inflammatory reactions would be enormous at this particular dosage. This would discourage the skilled artisan to pursue a medicament using MDP.

Finally, in Masihi et al., the cell lines in which the muramyl peptides were tested for inhibition of HIV-1 are T-Tropic HIV-1 strains. Masihi et al. is silent with respect to the testing of these compounds in M-Tropic HIV-1 strains which clearly distinguishes the presently claimed invention from this reference, as discussed above.

In other words, Masihi et al. teach that MDP can inhibit HIV-1 infection in the late stages of the disease. Masihi et al. does not disclose nor demonstrate that MDP or any other muramyl peptide for that matter can target the early stages of HIV-1 infection, which is the most important stage to target.

It should be clear that silence in a reference is not a proper basis to maintain an obviousness rejection.

From the foregoing, favorable action in the form of a Notice of Allowance is respectfully requested and earnestly solicited.

If the Examiner has any questions concerning this application, he is requested to contact the undersigned at (703) 205-8000 in the Washington, D.C. area.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees

Serial Number: 08/809,650

required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17;  
particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By: ~~Margaret A. Gunning (Reg. No. 28,977)~~  
Gerald M. Murphy, Jr.  
Reg. No. 28,977

P. O. Box 747  
Falls Church, VA 22040-0747  
(703) 205-8000

GMM/MAA/bsh

Oren Cohen  
Audrey Kuster  
Anthony S. Fauci

# Host factors in the pathogenesis of HIV disease

Annex I

Authors' address:  
Oren J. Cohen, Audrey Kuster, Anthony S. Fauci  
National Institute of Allergy and Infectious  
Diseases, Laboratory of Immunoregulation,  
Bethesda, Maryland, USA

Correspondence to:  
Oren J. Cohen  
National Institute of Allergy and Infectious  
Diseases  
Laboratory of Immunoregulation  
10 Center Drive, MSC 1576  
Building 10 Room 15B11  
Bethesda MD 20892-1576  
USA  
Fax: +1 301 402 0070  
e-mail: OCohen@nida.gov

**Summary.** Host factors play an important role in determining rates of disease progression in human immunodeficiency virus (HIV)-infected individuals. HIV is able to subvert the host immune system by infecting CD4<sup>+</sup> T cells that normally orchestrate immune responses and by inducing the secretion of proinflammatory cytokines that the virus can utilize to its own replicative advantage. The recognition that certain chemokine receptors serve as necessary co-factors for HIV entry into its target cells as well as the fact that ligands for these receptors can modulate the efficiency of HIV infection has expanded the number and scope of host factors that may impact the pathogenesis of HIV disease. This area of investigation will no doubt yield novel therapeutic strategies for intervention in HIV disease, however, caution is warranted in light of the enormous complexity of the pleiotropic cytokine and chemokine networks and the uncertainty inherent in manipulating these systems.

HIV-infected long-term non-progressors represent an excellent model to study potential host factors involved in HIV disease pathogenesis. Genetic factors certainly have a major impact on the immune responses mounted by the host. In this regard, a polymorphism in the gene for the HIV co-receptor CC chemokine receptor 5 (CCR5), which serves as a co-receptor for macrophage (M)-tropic strains of HIV, affords a high degree of protection against HIV infection in individuals homozygous for the genetic defect and some degree of protection against disease progression in HIV-infected heterozygotes. HIV specific immune responses, including cytotoxic T lymphocyte (CTL) responses and humoralizing antibody responses, also appear to play salutary roles in protecting against disease progression.

## Introduction

The pathogenesis of human immunodeficiency virus (HIV) disease is complex and influenced by both viral and host factors (1). The multifactorial nature of HIV disease pathogenesis is reflected by the highly variable rates of disease progression that are observed in individuals infected with HIV. The importance of host factors in modulating rates of disease progression is further underscored by the observation that even individuals who were apparently infected from a common source had experienced widely variable clinical outcomes (2). A great deal of attention has been focused recently on the discovery of polymorphisms in the *cckr5* gene that have been gained by mutation in the transmembrane domain containing potential viral binding sites (3,4). These studies

Received 15 January 2000  
Accepted 15 February 2000  
© 2000 Blackwell Science Ltd  
Journal of Internal Medicine 247: 11-21  
Correspondence: Oren J. Cohen, National Institute of Allergy and Infectious Diseases, Laboratory of Immunoregulation, 10 Center Drive, MSC 1576, Building 10 Room 15B11, Bethesda, MD 20892-1576, USA.  
Fax: +1 301 402 0070.  
e-mail: OCohen@nida.gov

has confirmed earlier work that demonstrated high levels of viral replication throughout the course of HIV infection (7-9) and have greatly expanded our understanding of the dynamics of HIV replication *in vivo*. The remarkable consistency in quantitative estimates of the rate of turnover of plasma virus raises important questions regarding the observed variability in rates of disease progression. In this regard, future studies will need to address whether the rate of viral turnover varies according to stage of disease or whether it is an intrinsic characteristic of HIV infection. In either case, it is necessary to invoke host factors in order to explain the great variability in rates of clinical disease progression.

A delicate balance among a wide array of host factors likely determines the net rate of viral replication in HIV-infected individuals. Subversion of the human immune system by HIV (i.e. infection of cells that are critical components of an intact immune system, induction of the secretion of proinflammatory cytokines, and utilization of these products of immune activation for the replicative advantage of the virus) usually tips the balance in favor of the virus. The recent discovery that certain chemokine receptors (for example, CC chemokine receptor (CCR)5, CXC chemokine receptor (CXCR)4, CCR3, CCR2b, STRL33, Bonzo, and BOB) are utilized by different strains of HIV as co-factors to gain entry into cells has greatly expanded the number of candidate host factors that may influence the pathogenesis of HIV disease (10-18). The ability of the chemokine ligands of these receptors to block HIV entry into target cells and thereby tip the balance of immune control over virus replication in favor of the host is a new concept in the field of HIV pathogenesis that has major implications for potential therapeutic intervention.

Genetic factors may determine the outcome of interactions between virus and host in several ways. First, the host's HIV-specific immune responses are constrained by the individual's major histocompatibility complex (MHC) alleles. In addition, the recently discovered genetic defect in the CCR5 gene has a major impact on susceptibility to HIV infection in individuals homozygous for the defect, and on disease progression in HIV-infected individuals heterozygous for the defect. HIV-specific cellular and humoral immune responses likely play an important role in the control of viral replication, although the precise correlates of protective immunity have not been established. However, recent studies have shown that qualitative as well as quantitative features of these immune responses may be important modulators of disease progression.

Appreciation of the role of host factors in the pathogenesis of HIV disease should lead to the design of novel therapeutic strategies. The goal of tipping the balance in favor of the host

over virus replication may appear to be simple, however the extraordinary complexity of manipulating host factors to this end is fraught with many potential complications. The latter point is highlighted by the negative outcomes of clinical trials for bacterial sepsis. In targeted molecules thought to be directly involved in the pathogenesis of sepsis (for example lipopolysaccharide, interleukin-1 $\beta$ , and tumor necrosis factor (TNF)- $\alpha$ ) (19). The need to consider therapeutic options in the context of a balance between pro- and anti-inflammatory mediators and the need to consider distal interactions in a complex pleiotropic cytokine network apply not only to sepsis (20), but to HIV disease as well.

#### Cytokines and HIV disease: dysregulation of cytokine production

A highly complex network of cytokines operates to regulate the immune system. This network is redundant and pleiotropic, and operates in an autocrine and paracrine manner to stimulate or suppress cellular proliferation and differentiation, and to modulate immune function (21). Chronic immune activation induced by HIV infection and associated opportunistic infections results in dysregulation of the cytokine network. Many of the observed alterations in cytokine production contribute to HIV pathogenesis by further stimulating viral replication, suppressing the ability of the immune system to mount a strong antiviral response, and inducing cytokine-mediated cytopathic effects (1, 22-24).

Similar to other chronic infections, HIV infection is associated with increased expression of proinflammatory cytokines, especially during the later stages of disease (25). High levels of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 are secreted by peripheral blood mononuclear cells (PBMC) (25-30) and macrophages (31-34) from HIV-infected subjects. TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 are also found at elevated levels in the serum (35-40), cerebrospinal fluid (41-43), and tissues (44-49). High levels of expression of these cytokines, as well as interferon (IFN)- $\gamma$  (46, 47, 50) and IL-3 (47, 50), are particularly evident in lymphoid tissue, a major site of HIV replication throughout the course of disease (8, 9, 51, 52). Chronically activated and expanded CD8<sup>+</sup> T cells (47, 52) and macrophages (22, 53) are thought to be major contributors to the elevated cytokine levels observed in HIV-infected subjects.

In addition to alterations in cytokine production and chronic immune activation, HIV-specific upregulation of certain cellular surface molecules, e.g., production of proinflammatory cytokines can be upregulated in HIV-infected monocytes (54-56), CD8<sup>+</sup> T cells (57), and macrophages (58) in

HIV infection in vitro after treatment with HIV proteins, such as envelope gp120 and gp124 (55-62).

Another major disruption in the cytokine pattern observed in HIV disease is a progressive loss in the ability to produce immunoregulatory cytokines, such as IL-2 and IL-12 (63-66). IL-2 and IL-12 are crucial for effective cell-mediated immune responses, as they stimulate proliferation and lytic activity of cytotoxic T lymphocytes (CTL) and natural killer (NK) cells. These cell-mediated immune effectors represent the primary mechanism whereby most viral infections are cleared. In addition, IL-12 is essential for stimulating the production of T helper (Th)1-type cytokines, including IL-2 and IFN- $\gamma$ , that favor the development of cell-mediated immune responses (67-69). While it is clear that the Th1 limb of cellular immune responses is impaired during the course of HIV infection (65, 70-73), controversy surrounds the proposed dominance of Th2-like responses (i.e. secretion of IL-4, IL-5, and IL-10) during progression of HIV disease. Clerici et al. showed that stimulated PBMC from HIV-infected patients exhibit a preferential Th2 pattern of cytokine secretion with disease progression (65, 70, 71, 74); however, other investigators have found a skewing of the cytokine secretion pattern of T cells from HIV-infected patients toward a Th0 state (i.e. secretion of cytokines characteristic of both Th1 and Th2 patterns) rather than toward a Th2 state (7, 72, 73). In either case, the finding that HIV replication is more efficient in Th0 compared to Th1 clones (72, 75) highlights the importance of impaired Th1 responses in the pathogenesis of HIV disease (76).

#### Effects of cytokines on HIV replication

The effects of cytokines on HIV replication were recognized in early studies wherein activated PBMC (77), macrophages (55), and B cells (79) were shown to produce soluble factors that could dramatically upregulate HIV expression in acutely and chronically infected cells of the lymphocyte and macrophage lineages. These observations led to the identification of numerous cytokines that can directly influence HIV replication in infected cells (22, 24-30) (Fig. 1).

Cytokines that have been reported to upregulate HIV replication *in vitro* include IL-1 $\beta$ , IL-2, IL-3, IL-6, IL-7 (81), IL-12 (82, 83), IL-15 (82, 84), TNF- $\alpha$ , TNF- $\beta$ , and the colony-stimulating factors (CSF) macrophage (M)-CSF and granulocyte macrophage (GM)-CSF (reviewed in (24)). IFN- $\alpha$ , IFN- $\beta$ , and IL-16 (85, 86) are primarily suppressors of HIV production, whereas other cytokines, such as IL-4 (87), IL-10 (88, 89), IL-13 (87), IFN- $\gamma$  and TGF- $\beta$ , reduce or enhance viral replication depending on the infected cell type and the culture condi-




Fig. 1. Endogenous cytokines regulate viral replication in CD4<sup>+</sup> T cells. Numerous cytokines, particularly the proinflammatory cytokines TNF- $\alpha$ , IL-1 $\beta$  and IL-6, strongly upregulate viral replication. TNF- $\beta$  and IL-1 $\beta$  downregulate viral replication in the case of IL-10; this effect is at least in part due to downregulation of proinflammatory cytokines. The proinflammatory cytokines which are secreted by a variety of cell types including CD4<sup>+</sup> and CD8<sup>+</sup> monocyte/macrophages, strongly stimulate infection by HIV-1, whereas IL-10 inhibits infection with HIV-1. (Adapted from (13)).

tions (22, 24, 30). Many cytokines, such as the interferons and TNF- $\alpha$ , can influence HIV replication in both T cells and macrophages, while others, such as M-CSF, are cell lineage-specific. The effects of a particular cytokine are often greatly influenced by the activity of other cytokines present in the microenvironment. In this regard, certain cytokines have been demonstrated to act in a synergistic (88, 90, 91) or in an antagonistic (91, 93) manner with other cytokines in regulating HIV replication. Finally, cytokines are pleiotropic and the overall effects of a particular cytokine on HIV replication often reflect the balance of both HIV-inducing and HIV-inhibiting activities.

Proinflammatory cytokines, particularly TNF- $\alpha$ , are considered the most potent HIV-inducing cytokines and their mechanism of action is relatively well understood. Both TNF- $\alpha$  and IL-1 $\beta$  activate the cellular transcription factor nuclear factor (NF) $\kappa$ B (94, 95), a strong inducer of HIV long terminal repeat (LTR)-mediated transcription. IL-6 alone appears to increase HIV expression primarily by a post-transcriptional mechanism; however, IL-6 can synergize with NF $\kappa$ B-inducing cytokines to enhance HIV transcription (90). The role of endogenous proinflammatory cytokines in the regulation of HIV replication has been demonstrated in several cellular systems *in vitro*. The production of HIV by macrophages or PBMC, stimulated with physiologic mixtures of proinflammatory cytokine production, such as bacterial endotoxin or IL-1, can be partially or nearly completely abrogated by the addition of anti-proinflammatory cytokines (25, 92), neutralizing antibodies to specific cytokines (96), such as IL-1, or mixtures of these cytokines (97). In macrophages, the virus suppressive activity of these cytokines

such as IL-10 and TGF- $\beta$  is attributable largely to their ability to inhibit the secretion or activity of HIV-inducing proinflammatory cytokines (92, 93, 96, 99). HIV production by infected T cells is sensitive to both the antiinflammatory and the antiproliferative activity of such cytokines (D. Collier & A.S. Fauci, unpublished data).

Although the role of proinflammatory and antiinflammatory cytokines in the regulation of HIV replication *in vivo* has not been demonstrated conclusively, several lines of evidence suggest that these cytokines may be involved in regulating viral production. Administration of pentoxifylline, an inhibitor of the secretion and activity of TNF, to HIV-infected individuals was found to reduce HIV viremia in concert with a reduction in plasma levels of TNF- $\alpha$  (100, 101). The role of proinflammatory cytokines in maintaining steady-state levels of HIV replication is suggested by the observation that *in vivo* infusion of a single bolus of IL-10 to HIV-infected subjects resulted in a rapid and modest, albeit transient, decrease in plasma viremia (D. Weissman & A.S. Fauci, unpublished data). The kinetics of HIV suppression *in vivo* correlated with a dramatic reduction in the ability of cells from these subjects to be induced *in vitro* to secrete TNF- $\alpha$  and IL-1 $\beta$ . Furthermore, IL-10 has been found to inhibit acute HIV infection in severe combined immunodeficiency (SCID) mice engrafted with human fetal thymus and liver (102). The ability of IL-10 to suppress T-cell activation and proliferation likely also plays a prominent role in its ability to suppress HIV replication *in vivo* (103–105).

In addition to the use of immunosuppressive cytokines which may depress HIV-inducing immune responses, cytokines which stimulate T cells or antigen-presenting cells have been administered to HIV-infected subjects for a number of years. The use of cytokine-based therapies aimed at immune reconstitution in HIV disease has expanded over the past several years, particularly with the development of potent antiretroviral therapies that limit the potential for cytokine-mediated increases in viral replication. In this regard, administration of IL-2 to asymptomatic HIV-infected subjects receiving concomitant antiretroviral therapy results in significant and sustained increases in CD4 $^{+}$  T-cell numbers with no long-term effects on viremia (106–107). Similar immune reconstitution therapies are being proposed for IL-1 $\beta$ , IL-15, and IL-15 (84, 108–110). New studies are continuing to expand the list of cytokines for use as potential immunotherapeutic agents. A particularly interesting cytokine-based immunotherapeutic approach is suggested by a recent report demonstrating that transfection of a CD4 $^{+}$  T cell line with DN Fc $\epsilon$  gene (the 133 amino acid form of IL-4 renders cells virtually resistant to IL-4 infection) (83). IL-4-mediated inhibition of HIV in this system appears

to be due to interference with viral transcription (85, 86). This effect may be due to the ability of IL-4 to suppress T-cell activation (116, 117). The combination of IL-2 and IL-16 is a particularly attractive option that may synergistically enhance the expansion of CD4 $^{+}$  T cells (N. Rautava, personal communication).

In addition to the known cytokines mentioned previously, several unidentified soluble factors have been demonstrated to exert dramatic HIV-modulating activity. Foremost among these is the elusive CD8 $^{+}$  cell-derived HIV suppressive factor(s). While lysis activity is an important component of CD8 $^{+}$  cell-mediated HIV suppression (118), cell-free supernatants from cultures of activated CD8 $^{+}$  cells and cell lines are able to dramatically inhibit HIV replication in both T cells and macrophages (119). CD8 antiviral factor (CAF), first described by Walker et al. (120–122), is non-cytotoxic, suppresses HIV replication in a non-MHC-restricted manner at the level of HIV LTR transcription (123–125) and lacks identity to known cytokines (126).

A distinct group of HIV-suppressive factors secreted by CD8 $^{+}$  T cells was identified by Cohen et al. (127). These investigators attributed the HIV-suppressive activity of CD8 $^{+}$  cells to the combined activities of certain chemottractant cytokines (i.e. chemokines), including macrophage inflammatory protein (MIP)-1 $\alpha$ , MIP-1 $\beta$ , and RANTES (Regulated upon Activation, Normal T Cell Expressed and Secreted). An unexplained finding in the study by Cohen et al. was that although the combination of the  $\beta$  chemokines MIP-1 $\alpha$ , MIP-1 $\beta$ , and RANTES potently suppressed the replication of several M-tropic HIV strains, they had virtually no effect on the replication of the T-cell line adapted (TCLA) strain, HIV-1 HIB. Soon after this report, Feng et al. described the seven transmembrane orphan receptor fusin, previously known as CXSTR and HUMSTR and currently designated CXCR4, as a coreceptor for T-cell (T)-tropic strains of HIV (128–130). In addition, three groups described a new chemokine receptor, CXCR5, which bound MIP-1 $\alpha$ , MIP-1 $\beta$ , and RANTES as its natural ligands (128–130). In light of the previous results of Cohen et al., the obvious question that arose was whether CXCR5 might function as a coreceptor for M-tropic strains of HIV. A series of papers from five different laboratories elegantly demonstrated this to be the case (131–135). Chemokine receptors that are involved in HIV co-receptors are apparently involved in the fusion process that occurs between viral and target membranes (136–137). The general picture that has emerged is that M-tropic strains of HIV utilize primarily the chemokine receptor CXCR5, and to a lesser extent CXCR4 for fusion and the fusion CXCR4 ligand MIP-1 $\alpha$ , MIP-1 $\beta$ , and RANTES as the primary CXCR5 ligands. HIV strains that contain T-tropism require MIP-1 $\alpha$  fused to

employ the  $\alpha$  chemokine receptor (CXCR4) an interaction that is blocked by the CXCR4 ligand stromal derived factor-1 (SDF-1). Many primary T-tropic HIV isolates exhibit a broad range of CCR usage, including CXCR4 and CCR5 (132, 133). The recent discoveries of other HIV co-receptors have already made obsolete the simplistic notion that CCR5 and CXCR4 are the only important co-receptors for M- and T-tropic strains of HIV, respectively (17, 18, 134).

Numerous cell types produce a variety of chemokines (135, 136), and modulation of the production of these factors may influence HIV replication in a strain-specific manner (Fig. 1). Therefore, the overall effect of immune activation and the secretion of proinflammatory or immunoregulatory cytokines on HIV replication must now be considered in the context of potential influences on chemokine production, chemokine co-receptor expression, and the predominant viral quasispecies that is replicating *in vivo*. Chemokine production, induced during inflammation, is enhanced by several cytokines, including TNF- $\alpha$ , IL-1 $\beta$ , and immunoregulatory cytokines, such as IL-7 and IL-15 (135, 137-139). Thus, in HIV-infected subjects in the early stages of disease, the ability of TNF- $\alpha$  to stimulate  $\beta$ -chemokine production and thereby suppress M-tropic entry may override its HIV-inducing effects; however, in individuals harboring predominantly T-tropic quasispecies in the later stages of HIV disease, only the HIV-inducing activity of TNF- $\alpha$  would be influential. In fact, TNF- $\alpha$ -mediated induction of  $\beta$ -chemokine secretion may actually enhance entry and replication of T-tropic strains of HIV (A. Kinter & A.S. Fauci, unpublished data) (Fig. 2).

Similarly, cytokines that modulate the expression of chemokine receptors would be expected to exert variable strain-dependent effects on HIV replication and spread. In this regard, IL-2 has been shown to upregulate the expression of the T-tropic co-receptor CCR5 (140).

The puzzling bottleneck in HIV transmission that so heavily favors emergence of M-tropic, non-syncytium-inducing (NSI) strains of virus in the new host (141, 142) may in part be due to the differential regulatory patterns of the relevant HIV co-receptors (140, 143). In this regard, CCR5 expression is predominantly seen in previously activated, memory T cells (i.e. CD45<sup>RO</sup>CD45<sup>RA</sup><sup>+</sup>CD45<sup>RO</sup><sup>+</sup>), whereas CXCR4 expression is seen in naive, inactivated cells (i.e. CD45<sup>RO</sup>CD45<sup>RA</sup><sup>+</sup>CD45<sup>RO</sup><sup>+</sup>). It is therefore plausible that the profound degree of immune activation that occurs during acute HIV infection may result in high expression of CCR5 and low expression of CXCR4. M. Gouws & A.S. Fauci, unpublished data. Similarly, co-infection with various other pathogens may differentially modulate expression of HIV co-receptors and thereby exert selective pres-




Fig. 2. Proinflammatory cytokines, such as TNF- $\alpha$ , have potentially dichotomous effects on HIV replication. During the early stages of HIV disease (left), M-tropic strains of the virus predominate. Although TNF- $\alpha$  can transcriptionally upregulate HIV expression in infected cells, at the same time it induces expression of the  $\beta$ -chemokines (RANTES, MIP-1 $\alpha$ , and MIP-1 $\beta$ ). These chemokines occupy the HIV co-receptor CCR5, blocking entry of HIV into target cells. In contrast, T-tropic strains of HIV may predominate in the late stages of HIV disease (right). In this situation, the induction of  $\beta$ -CC by TNF- $\alpha$  cannot block T-tropic HIV entry via CXCR4, and in fact may enhance replication of T-tropic strains of HIV.

sure on HIV strains that use the co-receptors in question (H. Mendenhall, M. Mendenhall, & A.S. Fauci, unpublished data).

The observation that the natural ligands of CC chemokine receptors that are utilized as HIV co-receptors act as potent inhibitors of viral entry has dramatically broadened the known spectrum of HIV-inhibitory cytokines and the mechanisms whereby they influence HIV replication. Therapeutic approaches currently being considered include administration of HIV-inhibitory chemokines, chemokine antagonists that occupy HIV co-receptors without transmitting a signal (144), and chemokines that are engineered to bind to HIV co-receptors noncovalently and remain in the extracellular space (145). The relative efficacy of these approaches will depend on the relative usage of particular co-receptors in the population of replicating HIV strains in a given patient and on whether  $\beta$ -chemokines themselves are immunomodulatory.

Journal of Acquired Immune Deficiency Syndromes • Volume 25, Number 2, February 1, 2000



PBMC from asymptomatic HIV-infected individuals harboring predominantly M-tropic HIV strains (145), but not in PBMC from individuals with more advanced disease harboring predominantly T-tropic HIV strains (146) (A. Kinter & A.S. Fauci, unpublished data). Similarly, HIV isolates obtained longitudinally from infected individuals with rapid disease progression exhibit reduced sensitivity to inhibition by  $\beta$  chemokines *in vitro* over time (146, 147).

A cautionary note about these potential therapies comes from the known association of the transition from M-tropic NSI to T-tropic/syncytium-inducing (SI) viruses with disease progression. The transition from an NSI to SI virus may occur by mutation of only a few amino acid residues predominantly in the envelope V3 loop (148-155). The HIV envelope V3 loop has also been shown to be a major determinant of co-receptor usage (156). Given the error rate of viral reverse transcriptase and the rapid dynamics of viral replication, mutations in the HIV envelope gene that encode SI strains must appear very early in disease; however, failure of such mutants to emerge until late in the disease process indicates a change in the selective advantage of such a mutation during the course of disease progression. Because SI variants are able to use a broader range of entry co-receptors (for example, CXCR4) compared with NSI viruses, it is possible that SI variants emerge in response to high levels of  $\beta$  chemokines that block cellular entry of viruses which utilize CCR5 (i.e. predominantly NSI viruses) (152, 146, 147). This potential effect of  $\beta$  chemokines should be investigated since the emergence of T-tropic HIV strains *in vivo* is associated with rapid CD4<sup>+</sup> T cell decline and disease progression (157). Further caution is warranted in light of potential dichotomous effects of the  $\beta$  chemokines on HIV replication in different cell types (119, 158). The situation *in vivo* is no doubt highly complex, and multiple host factors as well as regulatory aspects of co-receptor expression in different tissue compartments likely determine the environment in which selection for NSI or SI variants is made (1, 140, 159).

While *in vitro* culture systems and cell line models have allowed investigators to identify numerous host factors that influence HIV replication and to delineate the mechanisms whereby these factors suppress or enhance viral replication, it is difficult to anticipate how manipulation of these factors will ultimately influence HIV replication *in vivo*. It is clear that host factors function within the context of an interactive immunoregulatory cytokine network and can have pleiotropic effects on HIV replication, some of which are viral strain-specific. Nevertheless, numerous host factors have proven or promising immunotherapeutic potential that should be further explored and pursued clinically for the treatment of HIV disease.

#### Immune activation

Within 6 months to 1 year following primary HIV infection, plasma viremia appears to stabilize to a steady state or set point that is a strong prognostic indicator of the rate of disease progression (60). Underlying this deceptively stable viremia is a high rate of virus production and clearance (approximately 10<sup>9</sup> virions/day) (61-69), which is produced by newly infected CD4<sup>+</sup> T-cells (60). Thus, even during clinically asymptomatic stages of HIV infection, persistent virus production serves as a potent source of immune activation and subsequent cytokine secretion; these activities, in turn, stimulate further viral replication.

*In vivo* cellular activation is essential for productive HIV infection of CD4<sup>+</sup> T cells (161, 162), and agents that interfere with T-cell activation dramatically inhibit HIV replication in these cells (163, 164). The role of immune activation in stimulating HIV replication *in vivo* is demonstrated by increases in viremia in HIV-infected individuals persistently or transiently exposed to exogenous immune stimuli. In this regard, HIV-infected natives of sub-Saharan Africa, who experience persistent immune activation due to chronic exposure to parasites and other pathogens, harbor high viral loads associated with rapid progression of HIV disease (165, 166). Similarly, coinfection with opportunistic pathogens, such as active tuberculosis (167-170) or pneumocystis pneumonia (171), results in dramatic increases in levels of plasma HIV viremia that return to baseline upon successful treatment of the opportunistic infection (OI). The source of elevated viremia during OI was suggested by a recent study demonstrating that lymphoid tissue macrophages produce high levels of HIV in the setting of OI (172).

Confirmation of the role of immune stimulation in HIV replication has been established in studies demonstrating that immunization of HIV-infected subjects with influenza (173) or tetanus toxoid (174) antigens results in transient but substantial increases in plasma viremia. Furthermore, PBMC from HIV-infected subjects were rendered more susceptible to HIV infection *in vitro* following immunization with tetanus toxoid (174).

#### Long-term non-progressors: a model to study host factors in the pathogenesis of HIV disease

In recent years, it has become clear that a small percentage of HIV-infected individuals can maintain stable viral progression and be free of disease for decades. These individuals are called long-term non-progressors (LTNP). The study of LTNP may provide a model to study host factors in the pathogenesis of HIV disease.

trary, however, a reasonable consensus definition includes documentation of HIV infection for more than 7 years, a CD4<sup>+</sup> T cell count greater than 600 cells/ $\mu$ L without significant decline over time, no symptoms of HIV induced disease, and no history of antiretroviral therapy (184). Although a minority of cases of long-term non progressive HIV infection may be associated with attenuated strains of HIV (185-189), most data suggest that viral attenuation is rare among long-term non-progressors, and that host factors play a dominant role in determining the state of non progression (189, 191, 190-192).

#### Genetic factors

Host genetic factors influence the rate of disease progression in HIV infection. A number of different mechanisms may be responsible for the observed associations between certain HLA haplotypes and different rates of HIV disease progression (3-96). The ability of certain HLA molecules to efficiently present immunodominant viral epitopes in order to generate cell-mediated immune responses may explain an association with slow disease progression. Conversely, other HLA molecules may promote immunopathogenic responses associated with more rapid disease progression. In a recent study, HLA-B27, B57, and B51 were most strongly associated with slow progression of HIV disease, while HLA-A23, B37, and B49 were associated with rapid progression (96). An HLA profile was developed that distinguished a 6-fold difference between rates of disease progression in rapid versus slow progressors. Other genetic factors linked to rates of HIV disease progression include allelic forms of the vitamin D-binding factor Gc (197), variant alleles of mannose-binding lectin (198), and the TNF  $\alpha$ 2 microsatellite allele (199).

CCR5 is a major co-receptor for M-tropic strains of HIV-1 (above) (10-14). A mutant allele of the CCR5 gene that contains an internal 32 base pair deletion resulting in a truncated protein (200-202) has a major impact on susceptibility to HIV infection and on rates of disease progression in HIV infected individuals. Homozygosity for the CCR5 mutation results in near-total protection from HIV-1 infection (200, 202-207). Heterozygosity for the CCR5 mutation results in decreased expression of CCR5 on the cell surface and reduced infectability of CD4<sup>+</sup> cells with M-tropic strains of HIV-1 compared to CD4<sup>+</sup> cells from CCR5 wild-type individuals (208). Although heterozygosity for CCR5 does not appear to afford protection against HIV-1 infection *in vivo*, it may confer partial protection against disease progression in HIV-infected individuals (159, 202-204, 209, 210). Protection against disease progression in CCR5 heterozygotes is due in part to the lower viral




Fig. 3. Levels of plasma viremia are indistinguishable among HIV-infected long-term non-progressors stratified by CCR5 genotype. Dark bars represent geometric means.

load "set point" after HIV seroconversion and a slower rate of CD4<sup>+</sup> T-cell depletion compared with CCR5 wild-type individuals (204).

Heterozygosity for the CCR5 mutation is significantly more common in cohorts of HIV-infected long-term non-progressors compared to HIV-infected control populations (159, 202, 209, 210). However, despite the fact that the frequency of CCR5 heterozygotes is increased 2 fold among non-progressors compared to HIV-infected controls, still fewer than 50% of non-progressors are CCR5 heterozygotes (202, 209). The possibility that CCR5 heterozygotes might constitute a subgroup among non-progressors with the lowest viral loads and most preserved CD4<sup>+</sup> T cell counts was investigated. Interestingly, CCR5 wild-type and heterozygous long-term non-progressors were indistinguishable with regard to multiple immunologic and virologic parameters of disease activity (209). Mean CD4<sup>+</sup> T-cell counts were 910 cells/ $\mu$ L among CCR5 wild-type non-progressors and 885 cells/ $\mu$ L among CCR5 heterozygous non-progressors. Geometric mean levels of plasma viremia were 2,104 HIV RNA copies/mL among CCR5 wild-type non-progressors and 1,995 HIV RNA copies/mL among CCR5 heterozygous non-progressors (209) (Fig. 3).

We have previously demonstrated that, in contrast to individuals with progressive disease, HIV-infected long-term non-progressors maintain intact lymphoid tissue architecture (186, 187). However, a great deal of heterogeneity among non-progressors is evident in the degree of to normal hyperplasia and viral trapping within germinal centers (186, 187). When stratified according to CCR5 genotype as wild-type and heterozygous non-progressors were again indistinguishable with regard to these parameters (Fig. 4).

2 A. 05  
10/1/00




Fig. 4. The degree of follicular hyperplasia and the degree of virus trapping in lymph node germinal centers are indistinguishable among HIV-infected long-term non-progressors stratified by CCR5 genotype. CD21 staining, which highlights the structure of the germinal centers, is present in the RNA detection of HIV in situ in all patients. Representative immunohistochemical micrographs. Representative lymph nodes from non-progressors with abundant and scant follicular hyperplasia and virus trapping are shown for each CCR5 genotype. (Adapted from [100].)

Taken together, these data indicate that although CCR5 heterozygotes have an increased chance of becoming non-progressors, HIV-infected CCR5 wild-type individuals may arrive at the same non-progressor phenotype by other mechanisms.

#### Host immune response CTL responses

HIV-specific CTL play an important role in the control, albeit incomplete, of HIV replication and spread [212, 213]. High precursor frequencies of HIV-specific CTL with broad specificity have been consistently detected in long-term non-progressors compared to progressors [180, 214–217]. Qualitative aspects of the HIV-specific CTL response are also important determinants of the efficacy of the CTL response in controlling viral replication. Maintenance of CTL responses specific for

viral core proteins is associated with a decreased risk of disease progression [217, 218]. This association does not appear to be true for CTL responses against other viral proteins. Recognition of a immunodominant CTL epitope presented by particular MHC class I alleles may result in potent anti-HIV activity [219] and may in part explain the association of certain MHC class I alleles with slower progression of HIV disease [194–196]. Furthermore, the skewing of the T-cell receptor V $\beta$  repertoire in HIV-infected patients has suggested that the ability to recruit an HIV-specific CTL response composed of a heterogeneous group of V $\beta$  families during primary infection is associated with better control of viral replication and a improved prognosis compared to mobilization and expansion of CTL from only one or two V $\beta$  families [220]. Control and maintenance of specialized CTL may be a common mechanism of viral clearance. This appears to be a strategy for virus persistence observed in certain strains

of lymphocyte chemokine receptors that rapidly and completely mobilize the host CTL response resulting in CTL exhaustion (i.e. high zone tolerance) (221). CTL exhaustion may occur to some degree in HIV infection where disappearance of certain originally expanded CTL clones can be demonstrated in the absence of viral escape mutations that might otherwise explain the phenomenon (222).

Taken together these observations argue against an immunopathogenic role for CTL in HIV disease (223) and in favor of a salutary role in the maintenance of low viral load and the state of non-progression. This inference is further supported by the demonstrated role of CTL in reducing levels of plasma viremia during primary HIV infection (224-225), and the association of progression to AIDS with late viral escape from a long-lived (9-12 years) immunodominant CTL response (226).

The host CTL response against HIV is constrained by the ability of the host's MHC class I alleles to bind to various viral epitopes, while the virus is constrained by the degree to which an escape mutation impairs viral fitness. These host-virus dynamics are extraordinarily complex given the large number of permutations of viral epitopes and MHC class I alleles. Viral mutations within CTL recognition epitopes (i.e. "escape mutants") are associated with increased levels of viral replication and progression of HIV disease (226-229). Viral escape mutants may thrive due to the release of CTL control over their replication and may also inhibit CTL responses against the pre-escape viral epitope (230, 231). However, certain viral escape mutations may be costly to viral fitness. In this regard, it has been reported that diffuse infiltrative CD8 lymphocytosis in HIV infection was associated with certain HLA types that apparently constrain evolution of viral sequence diversity in the envelope V3 loop (232). Other studies have highlighted the constraints on the host CTL response imposed by MHC class I alleles. It has been reported that in an HIV-infected individual, CTL clones specific for an HLA-B\*44 restricted epitope of gp120 displayed very limited diversity of T-cell receptor utilization (233). Furthermore limited plasticity of certain CTL responses in individuals with viral escape mutants, where the dominant CTL response may remain largely directed at the pre-escape viral epitope, has been demonstrated (234-235). The possibility that increased plasticity in the CTL response may allow the host to maintain more continuous and effective control over viral replication was suggested by studies demonstrating increased viral sequence diversity and generation of vigorous escape mutant-specific CTL responses in slow progressors (236-237). A mathematical model of CTL virus dynamics has been proposed that describes late progression as a result of viral sequence variation that escapes an immunodominant CTL

response and shifts the host response towards a weaker epitope (238). Thus, disease progression may be the result of fitness of viral escape mutants outpacing the plasticity of the host CTL response and slow progression may be the result of CTL plasticity overpowering viral escape mutants with limited fitness (239).

#### 4.2.2. HIV Infection Stage

Antiviral soluble factors elaborated by CD8<sup>+</sup> T cells may also play a role in non-progression of HIV infection. CAE, first described by Walker et al. (120-121), is non-cytolytic and non-MHC-restricted, inhibits viral replication at the level of HIV RNA transcription (123-125), and lacks identity to known cytokines (126). CAE activity was found to correlate with stage of disease (239). Fewer CD8<sup>+</sup> T cells from asymptomatic patients without significant CD4<sup>+</sup> T cell depletion were required to suppress viral replication *in vitro* compared to CD8<sup>+</sup> T cells from patients with advanced stage HIV disease. Studies of long term non-progressors have demonstrated more potent CD8<sup>+</sup> T cell derived soluble antiviral responses compared with progressors (131, 240).

RANTES, MIP-1 $\alpha$ , and MIP-1 $\beta$  are also important antiviral soluble factors secreted by CD8<sup>+</sup> T cells as well as other cell types (127, 145). These chemokines are natural ligands for the chemokine receptor and X-tropic HIV co-receptor CXCR5 and inhibit viral replication primarily at the level of entry. Conflicting data have been obtained regarding a relationship between levels of these chemokines and progression of HIV disease (132, 240-245). These conflicting data are not surprising since they came from studies of sera or stimulated PBMC. A recent report does, however, support a possible role for the CXCR5 chemokines in the protection of some exposed uninfected individuals against HIV infection. Upon stimulation with HIV antigen, CD4<sup>+</sup> T cells from these individuals secreted high levels of these chemokines that were capable of inhibiting the replication of X-tropic strains of HIV *in vitro* (246). Ultimately, levels of expression of these chemokines in lymphoid tissue, the primary site of HIV replication *in vivo*, may be the most relevant measurement with regard to an association with rates of disease progression.

CD8<sup>+</sup> T cell differential chemotaxis to chemokines (RANTES) against CXCR5 which attracts memory T cells (247) and SDF-1, the ligand for CXCR4 which attracts naive T cells (248) may influence the migration of T cells into and out of the effector site as well as potential CTL target cell recruitment. Results concerning the potential role of chemokines in HIV disease are conflicting and their effects on the host-virus interaction in viral replication, the maintenance of low levels of HIV



of neutralizing antibodies (185, 187, 253). It has been reported that the presence of HIV-1B neutralizing antibodies correlated with a more favorable prognosis (254). Subsequent studies demonstrated that the presence of neutralizing antibodies to primary HIV isolates and to autologous virus was associated with non-progression (255, 256). Furthermore, viral escape from neutralizing antibody responses is associated with emergence of the SI phenotype of HIV and with disease progression (255, 257). HIV-infected long-term non-progressors tend to maintain antibody responses that can neutralize a broad panel of primary isolates and also maintain neutralizing antibodies against autologous virus isolates; however, non-progressors are a heterogeneous group with regard to these neutralizing antibody responses (253, 255). Whether the maintenance of neutralizing antibodies in non-progressors is simply a marker for a relatively intact immune system or whether these antibodies play an active role in determining the state of non-progression remains unclear.

#### Lymphoid tissue: substrate for immune competence

The morphologic abnormalities of lymphoid tissue associated with HIV disease progression are important determinants of immunodeficiency (8, 258-264). Despite the long period of HIV infection in long-term non-progressors, histopathologic

examination of lymph node biopsies from these individuals revealed only mild HIV-related abnormalities such as follicular hyperplasia (180, 211), follicular involution, fibrosis, and lymphocyte depletion, associated with progressive HIV disease were found to be lacking in lymph nodes from non-progressors. The degree of follicular hyperplasia seen in non-progressors is significantly lower compared to that seen in progressors, and quantitative estimates would show evidence of atrophic germinal centers extending into the nodal medulla (180, 211). It is likely that preservation of lymphoid architecture in non-progressors is a reflection of the lower levels of viral replication over time in these individuals. Regardless of the mechanisms responsible for lower levels of viral replication in non-progressors, preservation of lymphoid tissue architecture is a critical component of the immunocompetence observed. This further highlights the need to understand the mechanisms responsible for the destruction of lymphoid architecture during progression of HIV disease. If immunorestorative strategies in advanced HIV infection are to be successful, substrate for the generation of immune responses (i.e. intact lymphoid tissue) must be present, necessitating the prevention or reversal of the histopathologic abnormalities of lymphoid tissue associated with HIV disease progression.

#### References

1. Fauci AS. Host factors and the pathogenesis of HIV-induced disease. *Nature* 1996; 381: 529-533.
2. Liu S-L, et al. Divergent patterns of progression to AIDS after infection from the same source. *Human Immunodeficiency Virus Type 1 Infection and Antiviral Responses*. *J Virol* 1997; 71: 1784-1795.
3. Ho OD, Neumann AU, Persson AS, Cohen W, Leonard JM, Nickbarg M. Rapid turnover of plasma virions and CD4 lymphocytes in HIV-1 infection. *Nature* 1995; 373: 1230-1234.
4. Wei X, et al. Viral dynamics in human immunodeficiency virus type-1 infection. *Nature* 1995; 373: 117-122.
5. Persson AS, Neumann AU, Nickbarg M, Leonard JM, Ho OD. HIV-1 dynamics in vivo: virion clearance rate, infected cell life span and viral generation time. *Science* 1996; 271: 1583-1586.
6. Persson AS, et al. Decay characteristics of HIV-1 infected compartments during combination therapy. *Nature* 1996; 381: 436-439.
7. Pridak M, et al. High levels of HIV-1 in plasma during all stages of infection determined by competitive ELISA. *Science* 1993; 259: 1749-1754.
8. Pantaleo G, et al. HIV infection, chronic and progressive lymphoid tissue during the clinically latent stage of disease. *Nature* 1993; 362: 355-358.
9. Enbomson E, et al. Massive covert infection of helper T lymphocytes and macrophages by HIV during the incubation period of AIDS. *Nature* 1993; 362: 359-361.
10. Deng M, et al. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 1996; 381: 661-666.
11. Dragic T, et al. HIV entry into CD4+ cells is mediated by the chemokine receptors CXCR4 and CXCR5. *Nature* 1996; 381: 667-671.
12. Okamoto C, et al. CXCR4 and CXCR5: VIP and MIP-12 receptors with distinct roles for chemokine signaling. *Science* 1996; 272: 1953-1956.
13. Choe H, et al. The chemokine receptors CXCR4 and CXCR5 facilitate infection by primary HIV-1 isolates. *Cell* 1998; 95: 1335-1349.
14. Doherty B, et al. A functional primary HIV-1 isolate uses fusin and the chemokine receptors CXCR4, CXCR5, and CXCR6 as fusion cofactors. *Cell* 1998; 95: 1351-1355.
15. Berg M, Boudreau N, Bergeron J. HIV-1 entry: a major function of DNA binding is to identify a membrane domain of a protein involved in entry. *Science* 1996; 272: 371-377.
16. Dermer C, et al. Doherty B, Bergeron J, Choe H. A chemokine receptor domain mediates HIV-1 infection of T cells. *Science* 1998; 280: 1584-1587.

1. Gaudin M, Alexander S, Finkel K, Shinnick G, Berger E, Barner J. SFRS33: a novel, chemokine receptor-like protein functions as a chemokine receptor for both macrophage tropic and T-cell-tropic HIV-1. *J Exp Med* 1997;185:1015-1023.
2. Deng H, Gnanapavan D, KewarRaman V, Gnanan D. Expression cloning of new receptors used by simian and human immunodeficiency viruses. *Nature* 1997;388:126-130.
3. Glauser M. The inflammatory cytokines: New developments in the pathophysiology and treatment of septic shock. *Drugs* 1996;52 (suppl 2):9-17.
4. Bone R. Immunologic dissonance: a continuing evolution in our understanding of the systemic inflammatory response syndrome (SIRS) and the multiple organ dysfunction syndrome (MODS). *Ann Intern Med* 1996;125:690-691.
5. Paul W, Seider R. Lymphocyte responses and cytokines. *Cell* 1994;76:241-251.
6. Aloisi K, Pontiggia P, Medenica R, Rizzo S. Cytokine patterns in adults with AIDS. *Immunol Invest* 1997;26:341-350.
7. Miedema F. Immunological abnormalities in the natural history of HIV infection: mechanisms and clinical relevance. *Immunologic Rev* 1992;3:173-193.
8. Potti G, Fauci A. Role of cytokines in the pathogenesis of human immunodeficiency virus infection. In: Aggarwal S, Puri R, eds. *Human cytokines: their role in disease and therapy*. Cambridge, MA: Blackwell Science, 1995. p. 421-49.
9. Lau A, Der S, Read S, Williams B. Regulation of tumor necrosis factor receptor expression by acid-labile interferon- $\alpha$  from AIDS sera. *AIDS Res Hum Retroviruses* 1997;7:545-552.
10. Rieux-Lombard P, Modoux C, Cruchaud A, Dayer J. Purified blood monocytes from HIV-1 infected patients produce high levels of TNF- $\alpha$  and IL-1. *Clin Immunol Immunopathol* 1989;50:37-38.
11. Houe D, Haque A, Watro P, Beaudeau C, Mouton Y, Caron A. Production of tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interleukin-1 (IL-1) in patients with AIDS. Enhanced levels of TNF- $\alpha$  is related to a higher cytotoxic activity. *Clin Exp Immunol* 1989;78:329-333.
12. Voth R, et al. Differential gene expression of IFN- $\alpha$  and tumor necrosis factor- $\alpha$  in peripheral blood mononuclear cells from patients with AIDS-related complex and AIDS. *J Immunol* 1993;144:979-985.
13. Vyakarnam S, et al. Serum production of tumor necrosis factor- $\alpha$  and beta and interferon- $\gamma$  in HIV-1 infected individuals. *Clin Exp Immunol* 1994;94:106-111.
14. Weiss L, Hartner-Cavanagh N, de M, Gnanan D, Karachmand N. HIV infection is associated with the increased production of interleukin-12 (IL-12) and reduced interleukin-10 (IL-10) alpha serum. *AIDS* 1997;11:69-72.
15. Muller A, Miller R, Enay N, Meager A, Semple S, Roelke T. Production of tumor necrosis factor- $\alpha$  by blood and lung mononuclear phagocytes from patients with human immunodeficiency virus related lung disease. *Am J Respir Cell Mol Biol* 1994;9:48-54.
16. Krishnan V, Meager A, Mitchell D, Enchong A. Alveolar macrophages in AIDS patients increased spontaneous tumor necrosis factor- $\alpha$  production in pneumocystis carinii pneumonia. *Clin Exp Immunol* 1990;80:156-161.
17. Antinori A, et al. Alveolar macrophages from AIDS patients spontaneously produce elevated levels of TNF- $\alpha$  in vitro. *Allergol Immunopathol* 1992;20:147-154.
18. Israel-Biet D, Cathelin J, Goldstein A, Anouilh J, Jeffrey A, Even P. Tumor necrosis factor production in HIV-seropositive subjects: relationship with lung opportunistic infections and HIV expression in alveolar macrophages. *J Immunol* 1991;147:470-474.
19. Roddy M, Sorrell S, Lange M, Grieco M. Tumor necrosis factor and HIV p24 antigen levels in serum of HIV-infected populations. *J Acquir Immune Defic Syndr* 1988;1:436-440.
20. Lohdevirta J, Maury C, Teppo K, Korpela H. Elevated levels of circulating cytokines, tumor necrosis factor- $\alpha$  in patients with acquired immunodeficiency syndrome. *Am J Med* 1988;85:137-139.
21. Arditi M, Kazar W, Tager A, Sordani R. Tumor necrosis factor- $\alpha$ , interleukin-1, and p24 antigen concentrations in the blood of various stages of human immunodeficiency virus-1 infection. *Arch Intern Med* 1990;150:445-449.
22. Sudo-Agata D, Vulliamy E, Marwede M, de Saint Martin J. Diagnostics Serum and IL-12 (IL-12) alpha, TNF- $\alpha$  and IL-10 receptor of IL-12 in HIV-1 infected individuals. *AIDS Res Hum Retroviruses* 1999;15:336-338.
23. Brown EC, et al. Infection with HIV-1 associate low levels of IL-10 in the serum of patients. *Immunol Lett* 1994;40:43-44.
24. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:45-46.
25. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:47-48.
26. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:49-50.
27. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:51-52.
28. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:53-54.
29. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:55-56.
30. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:57-58.
31. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:59-60.
32. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:61-62.
33. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:63-64.
34. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:65-66.
35. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:67-68.
36. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:69-70.
37. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:71-72.
38. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:73-74.
39. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:75-76.
40. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:77-78.
41. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:79-80.
42. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:81-82.
43. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:83-84.
44. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:85-86.
45. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:87-88.
46. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:89-90.
47. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:91-92.
48. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:93-94.
49. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:95-96.
50. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:97-98.
51. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:99-100.
52. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:101-102.
53. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:103-104.
54. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:105-106.
55. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:107-108.
56. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:109-110.
57. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:111-112.
58. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:113-114.
59. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:115-116.
60. Rautavaara J, et al. Interleukin-10 concentrations are elevated in HIV-1 infected individuals. *Immunol Lett* 1994;40:117-118.

43. Jansen AM, et al. Human immunodeficiency virus type 1 (HIV-1) glycoprotein 120 promotes cellular gamma interferon, tumor necrosis factor alpha (TNF-alpha), and TNF-beta when they encounter their target antigens. *J Virol* 1993; 67: 2844-2852.
44. Bover R, et al. Secretory repression of HIV infection: human monocytes/macrophages. *Pathobiology* 1994; 59: 119-122.
45. Chouksey K, Robbins PD, Fennie B, et al. Fautz AS. Viral antigen stimulation of the production of human monokines capable of regulating HIV-1 expression. *J Immunol* 1989; 143: 170-175.
46. Molina R, Sauter D, Byrn R, Dikatchis J, Gersmann J. Production of tumor necrosis factor alpha and interleukin-1 beta by monocyte cells infected with human immunodeficiency virus. *J Clin Invest* 1989; 84: 733-737.
47. Rieckmann P, Poli G, Fox C, Kehrl J, Fautz AS. Recombinant gp120 specifically enhances tumor necrosis factor-alpha production and IgA secretion in B lymphocytes from HIV-infected individuals but not from seronegative donors. *J Immunol* 1991; 147: 2922-2927.
48. Merrill J, Kiyonagi Y, Chen L. Interleukin-1 and tumor necrosis factor alpha can be induced from mononuclear phagocytes by human immunodeficiency virus type 1 binding to the CD4 receptor. *J Virol* 1989; 63: 4404-4408.
49. Brady H, Abraham D, Pennington D, Miles C, Jenkins S, Dierckx E. Altered cytokine expression in T lymphocytes from human immunodeficiency virus T4 transgenic mice. *J Virol* 1995; 69: 7622-7629.
50. Hofman F, Wright A, Dehainwala M, Wong-Sealy F, Walker S. Exogenous viral protein activates human endothelial cells. *Blood* 1993; 82: 2774-2780.
51. Blazevic V, Heino M, Lagerstedt A, Rantala K, Krohn K. Interleukin-10 gene expression induced by HIV-1 for and Rev in the cells. *Acquir Immune Defic Syndr Hum Retrovir* 1996; 13: 228-234.
52. Tying S, et al. Synthetic peptides corresponding to sequences in HIV-1 envelope gp120 and gp120 enhance in vitro production of interleukin-1 and tumor necrosis factor but depress production of interleukin-6, interferon-gamma and interleukin-12. *J Virol Immunol* 1991; 4: 33-42.
53. Lane HC, Depper JM, Greene WL, Waldman TA, Fautz AS. Quantitative analysis of immune function in patients with HIV-1. Acquired immunodeficiency syndrome. Evidence for a selective defect in viral antigen recognition. *N Engl J Med* 1985; 313: 10-15.
54. Schenk RD, Clerici M, Dierckx E, Shearer GM. Limiting infection analysis of interleukin-2-producing T cells responsive to recall and alloantigens in human immunodeficiency virus-infected and uninfected individuals. *Int J Immunol* 1993; 23: 1-12.
55. Clerici M, et al. Restriction of HIV-1 infection-mediated immune responses by interleukin-12 in vitro. *Science* 1993; 262: 1711-1714.
56. Meynard L, Schuitemaker H. Altered T-cell dysfunction in HIV infection: analogy due to defective antigen-presenting cell function? *Immunol Today* 1993; 14: 15-18.
57. Ronchiani S, Human TH1 and TH2 subset regulation of differentiation and function: protection and immunopathology. *Int Arch Allergy Immunol* 1993; 99: 279-295.
58. Hsien CS, Macaronis SE, Tripp CS, White O'Garra A, Murphy KM. Development of TH1 CD4+ T cells through IL-12 produced by Listeria-induced macrophages. *Science* 1993; 260: 547-549.
59. Trinchieri G. Interleukin-2 and its role in the generation of TH1 cells. *Immunol Today* 1993; 14: 133-138.
60. Clerici M, et al. Changes in interleukin-1 and interleukin-4 production in asymptomatic human immunodeficiency virus-seropositive individuals. *J Clin Invest* 1993; 91: 759-765.
61. Clerici M, et al. Role of interleukin-12 in T helper cell dysfunction in asymptomatic individuals infected with the human immunodeficiency virus. *J Clin Invest* 1994; 93: 768-775.
62. Maggi E, et al. Ability of HIV to promote TH1 to TH0 shift and to replicate preferentially in TH1 and TH0 cells. *Science* 1994; 265: 244-248.
63. Meynard L, O'Garra A, Keen L, Schuitemaker H, Miedema F. Changes in cytokine secretion patterns of CD4+ T-cell clones in human immunodeficiency virus infection. *Blood* 1994; 84: 4262-4268.
64. Clerici M, Shearer GM. A TH1-TH2 switch: a critical step in the ecology of HIV infection. *Immunol Today* 1993; 14: 10-13.
65. Vyakarnam A, Maier P, Martin S, Wiggall H. TH1 cells specific for HIV-1 gag p24 are less efficient than TH0 cells in supporting HIV-1 replication and immune virus replication in TH0 cells. *Immunology* 1993; 86: 33-45.
66. Clerici M, et al. Type 1 cytokine production and low prevalence of viral infection in patients with long-term non-progressing HIV-1 infection. *AIDS Res Hum Retrovir* 1995; 11: 651-655.
67. Ficks TH, et al. Kinter AL, Dierckx EA, Fautz AS. Cellular regulation of expression of HIV-1 in a chronically infected promonocytic cell line. *Science* 1987; 236: 116-119.
68. Glavin K, et al. Molecular regulation of human monocyte-derived virus expression: the role of cellular transcription factors. *Immunol* 1993; 142: 1-13.
69. Rieckmann P, et al. Fautz AS. Activated T lymphocytes from human immunodeficiency virus-infected individuals induce virus expression in infected T cells and a promonocytic cell line. *J Exp Med* 1991; 173: 1-5.
70. Poli G, et al. The immunomodulation of HIV expression. *Semin Immunol* 1993; 3: 143-152.
71. Smithgall JE, Wong G, Trinchieri G, Hutter D. IL-12 up-regulates HIV-1 transcription in naturally infected promonocytic mononuclear cells. *J Immunol* 1996; 156: 1374-1380.
72. Bayard M, Nedjmi M, Hsien H, Hsien H, Hsien W, Hsien J. Different effects of interleukin-12, interleukin-10, and interleukin-13 on human immunodeficiency virus type 1 transcription in vitro. *Clin Diagn Lab Immunol* 1999; 3: 107-111.
73. Kinter AL, Boudreau SV, Hsien H, Jackson R, Fautz AS. Interleukin-2 induces CD4+ T cell mediated suppression of human immunodeficiency virus replication in CD4+ T cells and blocks follicular dendritic cell mediated virus expression. *Proc Natl Acad Sci USA* 1992; 89: 10985-10989.
74. Lacey D, et al. In vivo immunologic and virologic effects of recombinant human peripheral blood mononuclear cell derived normal donor human immunodeficiency virus type 1 infected patients. *Clin Diagn Lab Immunol* 1999; 6: 100-104.
75. Zhou P, Goulet S, Lefebvre C, Lefebvre J, Norkko A. Human CD4+ T cells infected with HIV-1 express a novel expression of interleukin-10. *Nat Med* 1996; 2: 103-106.
76. Masiamek A, Parada N, Chakrabarti S, Center D, Kornienko H. HIV-1 infection suppresses HIV-1 promoter activity. *J Immunol* 1997; 158: 100-104.
77. Nair H, et al. HIV-1 infection suppresses the expression of interleukin-12 in monocytes. *Immunol* 1996; 142: 100-104.



- 88 Weissman D, Poli G, Fauci AS. IL-10 synergizes with multiple cytokines in enhancing HIV production in cells of monocytic lineage. *J Acquir Immune Defic Syndr Hum Retrovir* 1995;9:442-449.
- 89 Finnegan A, et al. IL-10 cooperates with TNF- $\alpha$  to activate HIV-1 from latency and actively infected cells of monocyte/macrophage lineage. *J Immunol* 1996;156:841-851.
- 90 Poli G, et al. Interleukin 6 induces human immunodeficiency virus expression in infected monocytic cells alone and in synergy with tumor necrosis factor alpha by transcriptional and post-transcriptional mechanisms. *J Exp Med* 1992;171:511-518.
- 91 Poli G, Kinter AL, Fauci AS. Interleukin 1 induces expression of the human immunodeficiency virus alone and in synergy with interleukin 6 in chronically infected U1 cells: inhibition of inductive effects by the interleukin 1 receptor antagonist. *Proc Natl Acad Sci USA* 1994;91:108-112.
- 92 Schuitemaker H, et al. Proliferation-dependent HIV-1 infection of monocytes occurs during differentiation into macrophages. *J Clin Invest* 1991;89:1154-1160.
- 93 Poli G, Kinter AL, Justement JS, Broder R, Kalish JL, Fauci AS. Transforming growth factor beta suppresses human immunodeficiency virus expression and replication in infected cells of the monocyte/macrophage lineage. *J Exp Med* 1991;173:589-597.
- 94 Okbom L, Kunkel S, Nabel GJ. Tumor necrosis factor alpha and interleukin 1 stimulate the human immunodeficiency virus enhancer by activation of the nuclear factor kappa B. *Proc Natl Acad Sci USA* 1989;86:2336-2340.
- 95 Dun EJ, Maury WJ, Hooks TM, Fauci AS, Raczyn AB. Tumor necrosis factor alpha activates human immunodeficiency virus type 1 through induction of nuclear factor binding to the NF-kappa B sites in the long terminal repeat. *Proc Natl Acad Sci USA* 1989;86:5971-5975.
- 96 Colella D, Kinter AL, Hardy JC, Poli G, Fauci AS. Modulation of endogenous IL-1 beta and IL-1 receptor antagonist results in opposing effects on HIV expression in chronically infected monocytic cells. *J Immunol* 1996;156:3501-3506.
- 97 Weissman D, Poli G, Fauci AS. Interleukin 10 blocks HIV replication in macrophages by inhibiting the autocrine loop of tumor necrosis factor alpha and interleukin 6 induction of virus. *AIDS Res Hum Retroviruses* 1994;10:1099-1106.
- 98 Kinter AL, Poli G, Hardy JC, Fauci AS. HIV replication in U1 macrophage peripheral blood mononuclear cells is driven in an autocrine-paracrine manner by endogenous cytokines. *J Immunol* 1993;151:2448-2454.
- 99 Schuitemaker H, et al. IL-10 is a potent inhibitor of HIV-1 replication in macrophages infected with HIV-1 virus. *Proc Natl Acad Sci USA* 1994;91:558-591.
- 100 Wachs PS, et al. Bone marrow biopsy in human immunodeficiency virus-seropositive persons with tuberculosis: a randomized controlled trial. *J Infect Dis* 1995;171:727-733.
- 101 Clerici M, et al. T-tropic line, monocytes, T-mediated immunity and reduced human immunodeficiency virus (HIV) plasma viremia in asymptomatic HIV-seropositive persons. *J Infect Dis* 1994;170:210-215.
- 102 Kollmann TR, et al. Inhibition of acute invasive human immunodeficiency virus infection by human interleukin 10 treatment of SCID mice implanted with human fetal thymus and liver. *Proc Natl Acad Sci USA* 1994;91:3126-3131.
- 103 Maschke R, et al. IL-10 inhibits HIV-1 replication and is induced by it. *Biochem Biophys Res Commun* 1994;202:374-383.
- 104 Moore KW, O'Garra A, de Waard-Marciniak E, Vieira R, Mosmann TR. Interleukin 10. *Annu Rev Immunol* 1993;11:155-190.
- 105 Ho A, Moore KW. Interleukin-10 and its receptor. *Ther Immunol* 1994;1:171-185.
- 106 Kovacs JA, et al. Controlled trial of interleukin-12 infusion in patients infected with the human immunodeficiency virus. *N Engl J Med* 1996;335:1150-1156.
- 107 Kinter AL, Fauci AS. Interleukin-2 and human immunodeficiency virus infection: pathogenesis, immunology and potential for immunologic enhancement. *Immunol Rev* 1995;151:1-15.
- 108 Seder R, Grabstein K, Berzonsky J, McDiver C. Cytokine interactions in human immunodeficiency virus infection: individuals infected with HIV-1 IL-2 and IL-15. *J Exp Med* 1995;181:1067-1077.
- 109 Kanatani T, Ando S, E. Taketomi Y, Levin N. IL-15 stimulates the expansion of AIDS virus-specific CTL. *J Immunol* 1994;152:363-368.
- 110 Smith R, Roberts R, Ark A, Nguyen L, Thomas F, Sautin F, Roman. Immunodeficiency virus (HIV) type 1 gp120 helps to cell-mediated cytotoxicity (CMC) and natural killer (NK) activity in HIV-infected cells. HIV-1 suppresses enhancement of NK activity by IL-12 and IL-15. *Immunopharmacol Immunopathol* 1997;20:1-11.
- 111 Williams JC, et al. IL-15 enhances human immunodeficiency virus infection. *Immunol* 1996;153:5978-5982.
- 112 Skrzewski K, Zarowski C. Interleukin 15: characterization and biological properties. *Immunol Rev* 1995;80:167-178.
- 113 Munder M, Gombosi S. Tumor necrosis factor- $\alpha$  modulates HIV-1 infection of macrophage-specific T-regulatory cells. *Rev Immunol* 1994;14:533-537.
- 114 Gately M, Murguev M. Interleukin-2 potentiates T cell applications in the treatment and prevention of infectious diseases. *Drugs* 1994;58:15S, suppl 2: 18-15.
- 115 Trinchieri G, Sironi P. The role of interleukin 12 in the immune response, disease and therapy. *Immunol Today* 1997;18:400-403.
- 116 Chackraborty A, et al. IL-15 inhibition of CD34 $^{+}$  T-dependent lymphocyte activation and proliferation. *J Immunol* 1996;157:5210-5216.
- 117 Thekkumkara V, Center D, Nicol J, Fine G, Kornfeld H, Chackraborty W. CD4 $^{+}$  and IL-15 inhibit the raten synapsis reaction. *J Immunol* 1996;157:1958-1964.
- 118 Yang J, et al. Suppression of human immunodeficiency virus type 1 replication by CD4 $^{+}$  cells: evidence for HLA class I-mediated triggering of cytotoxic and noncytotoxic mediated virus. *Virology* 1997;231:11-18.
- 119 Munder M, Gombosi S. Combining IL-12 with IL-15: a CD4 $^{+}$  T-cell-mediated anti-infectious virus that chemokines (CXCR4, MIP-1 $\alpha$  and MIP-1 $\beta$ ) suppress viral replication in macrophages. *Proc Natl Acad Sci USA* 1996;93:1097-1101.
- 120 Walker AS, Moore KW, Sacks DP, Sacks DA. CD4 $^{+}$  T lymphocytes control HIV infection in the absence of virus replication. *Science* 1996;274:1161-1164.
- 121 Walker M, Moore KW. Antiviral lymphokine produced by CD4 $^{+}$  T lymphocytes suppresses HIV replication. *Immunology* 1989;66:614-619.
- 122 Walker M, Frankham A, Brown B, Sacks DA. CD4 $^{+}$  T lymphocytes suppress immunodeficiency virus infection in the absence of CD4 $^{+}$  T lymphocytes. *Immunology* 1990;69:1-7.

11. Choe KH, Venzon RD, Baruch A, et al. Suppression of HIV-1 lymphocyte-mediated inhibition of HIV-1 long terminal repeat transcription by a novel antiviral mechanism. *AIDS Res Hum Retroviruses* 1993; **9**: 1079-1086.
12. Cope land KF, McKay PL, Rotherham KL. Suppression of activation of the human immunodeficiency virus long terminal repeat by CD8+ T cells: chemokines are specific. *AIDS Res Hum Retroviruses* 1995; **11**: 1371-1376.
13. Mackewicz C, Backbourn DJ, Levy JA. CD8+ T cells suppress human immunodeficiency virus replication by inhibiting viral transcription. *Proc Natl Acad Sci USA* 1995; **92**: 1305-1312.
14. Mackewicz CE, Orlando H, Levy JA. Effect of cytokines on HIV replication in CD4+ lymphocytes: lack of identity with the CD8+ cell antiviral factor. *Cell Immunol* 1994; **151**: 129-143.
15. Fu R, DeVico A, Garzino-Demo A, Anya S, Gallo R, Lusso P. Identification of RANTES, MIP-1 $\alpha$ , and MIP-1 $\beta$  as the major HIV suppressive factors produced by CD8+ T cells. *Science* 1995; **270**: 1817-1821.
16. Samson M, Labbe O, Mullereau C, Vassart G, Parmentier M. Molecular cloning and functional expression of a new CC chemokine receptor gene, CC-CCR5. *Biochemistry* 1995; **35**: 3367-3369.
17. Combadiere C, Ahuja S, Follis H, Murphy P. Cloning and functional expression of CC-CCR5, a human monocyte CC chemokine receptor selective for MIP-1 $\alpha$  (alpha), MIP-1 $\beta$  (beta), and RANTES. *Leukoc Biol* 1996; **60**: 147-157.
18. Raport C, Gao X, Schwab R, Gray P, Charo I. Molecular cloning and functional characterization of a novel human CC chemokine receptor (CCR5) for RANTES, MIP-1 $\beta$ , and MIP-1 $\alpha$ . *J Biol Chem* 1996; **271**: 1716-1720.
19. Oraveer T, Pail M, Norcross M, et al. Chemokine inhibition of monocytotropic HIV-1 infection: interference with virus binding fusion step. *J Immunol* 1996; **157**: 1179-1187.
20. Simmons G, et al. Primary viremia inducing human immunodeficiency virus type-1 isolates are dual-tropic and most can use either CXCR4 or CCR5 as coreceptors for virus entry. *J Virol* 1996; **70**: 8355-8360.
21. Zhang L, Huang Y, He T, Cao Y, et al. HIV subtype and second receptor use. *Nature* 1996; **383**: 768.
22. He T, et al. CCR5 and CXCR4 are coreceptors for HIV-1 infection in humans. *Nature* 1996; **383**: 90-93.
23. Schall T. Biology of the RANTES/lymphotoxin family. *Cytokines* 1991; **3**: 165-183.
24. Baggiolini M, Dewald B, Moser B. Human chemokines: an update. *Annu Rev Immunol* 1997; **15**: 675-713.
25. Nelson P, Kim H, Manning W, Baruch A, Krensky V. Genomic organization and transcriptional regulation of the RANTES chemokine gene. *Immunol* 1993; **151**: 260-267.
26. Rabinowitz P, Rabinowitz M, Sedick M, Schall T, McCull S. Expression of the chemokine RANTES in human rheumatoid synovial fibroblasts: Differential regulation of RANTES and interleukin-8 genes by inflammatory cytokines. *J Biol Chem* 1992; **268**: 5834-5837.
27. Schall T, Bacon N. Chemokines, cytokines, trafficking, and inflammation. *Curr Opin Immunol* 1993; **5**: 365-373.
28. Diehl C, Wu L, Huxley J, Springer T, Mackay L. The HIV coreceptors CXCR4 and CCR5 are differentially expressed and regulated in human T lymphocytes. *Proc Natl Acad Sci USA* 1997; **94**: 1925-1930.
29. Zhu T, et al. Genotypic and phenotypic characterization of HIV-1 in patients with primary infection. *Science* 1993; **261**: 1839-1841.
30. Zhang LQ, Mackenzie P, Cleland A, Holmes EC, Leigh-Brown AJ, Simmons P. Selection for specific sequences in the external envelope protein of human immunodeficiency virus type-1 upon primary infection. *J Virol* 1993; **67**: 3345-3356.
31. Unutmaz D, Littman D. Expression pattern of HIV-1 coreceptors on T cells: implications for viral transmission and lymphocyte homeostasis. *Proc Natl Acad Sci USA* 1997; **94**: 3163-3168.
32. Simmons G, et al. Potent inhibition of HIV-1 infectivity in macrophages and lymphocytes by a novel CCR5 antagonist. *Science* 1997; **276**: 276-279.
33. Kinter AL, et al. HIV replication in CD4+ T cells of HIV-infected and healthy is regulated by a balance between the viral suppressive effects of endogenous beta chemokines and the virus-inhibitory effects of other endogenous cytokines. *Proc Natl Acad Sci USA* 1996; **93**: 14076-14081.
34. Jansson M, et al. Sensitivity to inhibition by beta chemokines correlates with biological phenotypes of primary HIV-1 isolates. *Proc Natl Acad Sci USA* 1996; **93**: 15382-15387.
35. Connor R, Sheridan AE, Ceradini D, Chiswick L, Landau N. Change in coreceptor use correlates with disease progression in HIV-1 infected individuals. *J Exp Med* 1997; **185**: 1061-1068.
36. Park Higgins D, Ching M, Mayer C, Bauer D, Levy JA, Chao D. Human immunodeficiency virus type-1 isolates of high range replication capability are selected by the envelope region of the viral genome. *J Virol* 1990; **64**: 4034-4039.
37. Jelling H, et al. Human immunodeficiency virus type-1 envelope structure for the envelope glycoprotein: effect on virus formation and infection. *Science* 1991; **256**: 1171-1175.
38. Kiken T, Jelling H, E. Kiklen W, Teramoto N, Iwamoto T. Evolution of the V3 envelope domain in proviral sequences and isolation of human immunodeficiency virus type-1 during transition of the viral biological phenotype. *Virology* 1992; **66**: 1611-1617.
39. Liang SS, Brice CJ, Levy JA. Clonal diversification of the envelope V3 loop is the primary determinant of infectivity in HIV-1. *Science* 1991; **253**: 71-74.
40. Takeuchi Y, Akase M, Murawski K, Shimizu N, Hoshino H. Host range mutant of human immunodeficiency virus type-1: introduction of cell tropism by a single point mutation at the neutralization epitope in the env gene. *J Virol* 1991; **65**: 170-173.
41. Cheng M, et al. Shida T, Levy JA. Host range, infectivity, and pathogenic properties of human immunodeficiency virus type-1 are determined by a very few amino acid changes in gp120. *J Virol* 1991; **65**: 6931-6937.
42. Huchard RA, et al. Phenotype-associated sequence variation in the third variable domain of the human immunodeficiency virus type-1 gp120 molecule. *J Virol* 1992; **66**: 1834-1837.
43. Bhattacharya D, Broder BR, Gulland L. Positioning of amino acid charged residues in the V3 loop correlates with HIV type-1 virus from distinct phenotypes. *AIDS Res Hum Retroviruses* 1996; **12**: 83-93.
44. Gao X, Fu R, DeVico A, Garzino-Demo A, Gallo R, Lusso P. The first domain of HIV-1 envelope glycoprotein gp120 is critical for chemokine-mediated block of infection. *Nat Med* 1996; **2**: 1144-1147.
45. Kim N, et al. Prognostic value of HIV-1 viremia in predicting phenotype for rate of CD4+ cell depletion and progression to AIDS. *Ann Intern Med* 1996; **125**: 631-638.
46. Schindelman-Ross J, Berman M, Bukrinsky M. Chemokines and HIV replication. *Nature* 1996; **383**: 18.
47. Michael N, et al. The effect of chemokine and CCR5 gene polymorphisms on HIV-1 transmission and virus progression in the blood. *Proc Natl Acad Sci USA* 1997; **94**: 1061-1066.

160. Mellors JW, et al. Quantitation of HIV-1 RNA in plasma predicts outcome after zidovudine therapy. *Ann Intern Med* 1995;122:573-579.
161. Zack JA, Arrighi SJ, Weissman SR, Gu AS, Haislip A, Chen JS. HIV-1 entry into quiescent primary lymphocytes: molecular analysis reveals a latent, latent viral structure. *Cell* 1990;61:213-222.
162. Bukrinsky M, Stanwick J, Dempsey M, Stevenson M. Quiescent T lymphocytes as an inducible virus reservoir in HIV-1 infection. *Science* 1991;254:473-477.
163. Finberg R, et al. Selective elimination of HIV-1-infected cells with an interleukin-2 receptor-specific cytotoxic. *Science* 1991;252:1703-1705.
164. Belfi K, Ramilo O, Valera F. Combined use of an immunotoxin and cyclosporine to prevent both activated and quiescent peripheral blood T cells from producing type 1 human immunodeficiency virus. *Proc Natl Acad Sci USA* 1993;90:1411-1415.
165. Quinn T, et al. Serologic and immunologic studies in patients with AIDS in North America and Africa: The potential role of infectious agents as cofactors in human immunodeficiency virus infection. *J Am Med Assoc* 1987;257:2617-2621.
166. Beniswich Z, Kainkovich A, Weissman Z. Immune activation is a dominant factor in the pathogenesis of African AIDS. *Immunol Today* 1995;16:187-191.
167. Pape JW, Jean SS, Ho JL, Haefner A, Johnson WD. Effect of isoniazid prophylaxis on incidence of active tuberculosis and progression of HIV infection. *Lancet* 1993;342:268-272.
168. Wallis R, et al. Influence of tuberculosis on human immunodeficiency virus (HIV-1): enhanced cytokine expression and enhanced beta 2-microglobulin in HIV-1-associated tuberculosis. *Infect Dis* 1993;167:33-48.
169. Whalen C, Horsburgh CR, Hout J, Jahatt S, Simberloff M, Elner J. Accelerated course of human immunodeficiency virus infection after tuberculosis. *Am J Respir Crit Care Med* 1995;151:129-135.
170. Geyrho D, et al. Effect of Mycobacterium tuberculosis on HIV replication: Role of immune activation. *Immunol* 1995;157:127-137.
171. Israeli-Betz D, Cadranet J, Even P. Human immunodeficiency virus production by alveolar macrophages is increased during Pneumocystis carinii pneumonia. *Am Rev Respir Dis* 1995;148:108-112.
172. Davidson F, Fox C, Wahl S. Macrophages as a source of HIV during opportunistic infections. *Science* 1991;256:477-480.
173. Shapiro S, et al. Activation of virus replication after vaccination of HIV-1 infected individuals. *J Exp Med* 1995;182:1221-1227.
174. Stanley S, et al. Effect of immunization with a common recall antigen on viral expression in patients infected with human immunodeficiency virus type 1. *N Engl J Med* 1996;334:1024-1027.
175. Cohen M, et al. Long-term human immunodeficiency virus infection in asymptomatic homosexual and bisexual men with normal CD4+ lymphocyte counts: immunologic and virologic characteristics. *J Infect Dis* 1994;169:959-966.
176. Sheppard HW, Lang W, Ascher MS, Vittinghoff E, W. Haisstein W. The characterization of non-progressive, long-term HIV-1 infection with stable CD4+ T-cell levels. *AIDS* 1993;7:1159-1166.
177. Buchbinder SP, Katz MH, Hessel NA, O'Malley PM, Holmberg SD. Long-term HIV-1 infection without immunologic progression. *AIDS* 1994;8:1123-1128.
178. Keel JR, et al. Characteristics of long-term asymptomatic infection with human immunodeficiency virus type 1 in men with normal and low CD4+ cell counts. *J Infect Dis* 1994;169:1236-1243.
179. Morpur A, et al. Long-term survivors with HIV-1 infection: incubation period and longitudinal patterns of CD4+ lymphocytes. *J Acquir Immune Defic Syndr Hum Retrovir* 1995;8:496-505.
180. Pantaleo G, et al. Studies in subjects with long-term nonprogressive human immunodeficiency virus infection. *N Engl J Med* 1995;332:109-116.
181. Cao Y, Qin L, Zhang L, Salter L, Ho DD. Virologic and immunologic characterization of long-term survivors of human immunodeficiency virus type 1 infection. *N Engl J Med* 1995;332:201-208.
182. Vicenzi E, et al. Homophilia and nonprogressing human immunodeficiency virus type 1 infection. *Blood* 1997;89:191-200.
183. Balotta C, et al. Comparative biological and molecular determinants in HIV type 1-infected, long-term nonprogressors and recently infected individuals. *AIDS Res Hum Retrovir* 1995;11:331-341.
184. Schrago L, Young L, Fowler M, Strickland J, Vermund S. Long-term survivors of HIV-1 infection: biological and clinical challenges. *AIDS* 1994;8(Suppl 3):S108.
185. Leamon J, et al. Long-term survivors of HIV-1 infection: does it make a difference? *Lancet* 1995;340:363-367.
186. Meyer R, et al. Genetic structure of an extended quasi-species of HIV-1 from a blood transfusion donor and recipients. *Science* 1995;270:988-991.
187. Kirschhoff E, Greenough JC, Bretler DB, Sullivan JL, Desrosiers RC. Brief report: absence of intact nef sequences in a long-term survivor with nonprogressive HIV-1 infection. *N Engl J Med* 1995;332:224-232.
188. Vignati A, et al. Persistence of integrated provirus in a human immunodeficiency virus type 1-infected asymptomatic individual. *J Virol* 1995;69:5743-5753.
189. Macher M, et al. Defective accessory genes in a human immunodeficiency virus type 1-infected long-term survivor lacking detectable virus. *J Virol* 1995;69:4118-4126.
190. Macher M, Chang G, O'Leary LA, Tseng CJ, Birk DL, Sheppard HW. Functional characterization of human immunodeficiency virus type 1 nef genes in patients with divergent rates of disease progression. *J Virol* 1995;69:6758-6769.
191. Huang Y, Zhang L, Ho DD. Characterization of nef sequences in long-term survivors of human immunodeficiency virus type 1 infection. *J Virol* 1995;69:93-100.
192. Huang Y, Zhang L, Ho D. Biological characterization of nef in long-term survivors of human immunodeficiency virus type 1 infection. *J Virol* 1995;69:3142-3146.
193. Kallow RA, et al. A CCR5-DRB HLA antigen combination associated with rapid decline of Thelper lymphocytes in HIV-1 infection. *Lancet* 1990;335:937-939.
194. Klein MR, et al. Associations between HLA frequencies and pathogenic features of human immunodeficiency virus type 1 infection in seroconverters from the Amsterdam Cohort of Homosexual Men. *Infect Dis* 1994;169:1269-1279.
195. Keel JR, Klein MR, Just J, Kallow RA. The role of host genetics in the natural history of HIV-1 infection: the needles in the haystack. *AIDS* 1995;19(Suppl 3):S59-S67.
196. Kallow RA, et al. Influence of combination of human major histocompatibility complex genes on the course of HIV-1 infection. *N Engl J Med* 1995;333:405-411.
197. Eley G, et al. Association of different alleles of the major histocompatibility complex class II genes to clinical manifestations of human immunodeficiency virus infection. *J Infect Dis* 1995;171:104-110.

98. Carlier B, et al. Susceptibility to HIV infection and progression of AIDS in relation to various alleles of mannose-binding lectin. *Lancet* 1997;349:236-240.
99. Klein SH, et al. The TNF- $\alpha$  promoter allele is associated with the rate of HIV disease progression. *AIDS* 1997;11:423-428.
100. Samson M, et al. Resistance to HIV-1 infection in a sexless individual bearing multiple alleles of the CCR5 structural gene receptor. *Nature* 1996;382:722-725.
101. Liu R, et al. Homozygous defect in HIV-1 coreceptor accounts for resistance of some multiply-exposed individuals to HIV-1 infection. *Cell* 1996;86:167-177.
102. Zimmerman R, et al. Inherited resistance to HIV-1 conferred by an inactivating mutation in CC chemokine receptor 5: studies in populations with contrasting clinical phenotypes, defined racial backgrounds and unrelated risks. *Mol Med* 1997;3:23-36.
103. Dean M, et al. Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CCR5 structural gene. *Science* 1996;273:1856-1862.
104. Huang Y, et al. The role of a mutant CCR5 allele in HIV-1 transmission and disease progression. *Nat Med* 1996;2:1740-1743.
105. Bitt R, French R, Young J, Benoit B, Stewart G. HIV-1 infection in an individual homozygous for the CCR5 deletion allele. *Nat Med* 1997;3:252-253.
106. O'Brien T, et al. HIV-1 infection in a man homozygous for CCR5 $\Delta$ 32. *Lancet* 1997;349:1219.
107. Theodorou I, Meyer L, Magierowski M, Karlama C, Rouzioux C. HIV-1 infection in an individual homozygous for CCR5 $\Delta$ 32. *Lancet* 1997;349:1219-1220.
108. Wu L, et al. CCR5 levels and expression patterns correlate with infectability by macrophage-tropic HIV-1, in vitro. *J Exp Med* 1997;185:1681-1692.
109. Hemon D, et al. Heterozygosity for a defective gene for CC chemokine receptor 5 is not the sole determinant for the immunologic and virologic phenotype of HIV-infected long-term nonprogressors. *J Clin Invest* 1997;100:581-589.
110. Eugén-Olier J, et al. Heterozygosity for a deletion in the CCR5 gene leads to prolonged AIDS-free survival and lower CD4-T-cell fall in a cohort of HIV seropositive individuals. *AIDS* 1997;11:395-400.
111. Pantano G, Vaccarezza M, Graziosi F, Clementi M, Fauci AS. Antiviral immunity in HIV-1 infected long-term nonprogressors: CCR5 polymorphism. *AIDS* 1996;10:1149-1154.
112. Golden E, Williams A, McMahon A. Activated T cells - protection from disease progression - protection from infection. *Immunol Lett* 1996;51:175-178.
113. Weiss C, Walker B. HIV type 1-specific cytotoxic T lymphocytes and the control of HIV-1 reactivation. *Springer Semin Immunopathol* 1997;18:335-354.
114. Tarter P, et al. Cytotoxic T lymphocytes in asymptomatic long-term nonprogressing HIV-1 infection: breadth and spectrum of the response and relation to viral viral quasispecies in a person with prolonged infection and low viral load. *J Immunol* 1996;156:2619-2625.
115. Hauer T, et al. Strong cytotoxic T-cell and weak humoral antibody responses in a subset of persons with stable nonprogressing HIV type 1 infection. *AIDS Res Hum Retroviruses* 1996;12:585-592.
116. Rimoldi C, et al. High levels of anti-human immunodeficiency virus type 1 (HIV-1) memory cytotoxic T lymphocyte activity and low viral load are associated with lack of disease in HIV-1-infected long-term nonprogressors. *J Virol* 1995;69:5818-5821.
117. Klein MR, et al. Kinetics of gag-specific cytotoxic T lymphocyte responses during the clinical course of HIV-1 infection: a longitudinal analysis of rapid progressors and long-term asymptomatics. *J Exp Med* 1995;181:1365-1372.
118. Rizzetto Y, et al. Gag-specific cytotoxic responses to HIV type 1 are associated with a decreased risk of progression to AIDS-related complex or AIDS. *AIDS Res Hum Retroviruses* 1995;11:303-307.
119. Cockerill PJ, et al. Novel cross-reactive conserved and immunodominant cytotoxic T lymphocyte epitopes in slow progressors in HIV type 1 infection. *AIDS Res Hum Retroviruses* 1995;11:1671-1678.
120. Pantano G, et al. The putative function of the primary immune response to HIV infection as a prognosticator of disease progression independent of the initial level of plasma viremia. *Proc Natl Acad Sci USA* 1997;94:134-138.
121. Cockerill PJ, Lechner M, French R, Cockerill RM. Virus persistence in high infected immunodeficient mice by extremely potent neutralizing effect of T cells. *Nature* 1993;362:693-697.
122. Pantano G, et al. Evidence for HIV-1 dysregulation in initially exposed subjects with CD4-T-cell counts ranging from 100-400 cells/mm<sup>3</sup>. *AIDS* 1996;10:1155-1160.
123. Cockerill PJ, French R, Lechner M, Lechner RM. Induced immunopathology versus direct cytotoxicity by virus: implications for HIV and AIDS. *Immunol Today* 1993;14:121-127.
124. Kaul RA, et al. The viral association of cellular immune responses with the initial control of viremia in primary human immunodeficiency virus type 1 infection. *J Virol* 1994;68:4650-4655.
125. Borrow P, Lewicki H, Hahn BH, Shaw GM, Johnson VA. Virus-specific CD4+ cytotoxic T-lymphocyte activity associated with control of viremia in primary human immunodeficiency virus type 1 infection. *J Virol* 1994;68:6103-6110.
126. Goulder PR, et al. Late escape from an immunodominant cytotoxic T lymphocyte response associated with progression to AIDS. *Nat Med* 1997;3:211-217.
127. Phillips RE, et al. Human immunodeficiency virus genetic variation that can evade cytotoxic T-cell recognition. *Nature* 1997;384:433-434.
128. Koenig S, et al. Transfer of HIV-1-specific cytotoxic T lymphocytes to an AIDS patient leads to selection for mutant HIV variants and subsequent disease progression. *Nat Med* 1995;1:330-336.
129. Borrow P, et al. Antiviral pressure exerted by HIV-1-specific cytotoxic T lymphocytes (CTLs) during primary infection demonstrated by rapid selection of CTL-escape virus. *Nat Med* 1997;3:205-211.
130. Kienerman R, et al. Cytotoxic T-cell activity antagonized by naturally occurring HIV-1 Gag variants. *Nature* 1994;369:403-407.
131. Meyer U, et al. Cytotoxic T lymphocytes are inhibited by viable HIV mutants. *Science* 1995;270:1360-1361.
132. Hesse S, Rove S, Sawyer E, Wüthricher H. Certain HLA-DRB and -DQB1 major histocompatibility complex class II alleles are associated with a CD8 lymphocyte non-response to human immunodeficiency virus type 1 character 2e HIV-1 lymphocyte response heterogeneity and early disease progression. *Proc Natl Acad Sci USA* 1994;91:4411-4415.
133. Nairat SA, et al. Long-term analysis of T-cell receptor (TCR) gene usage by human immunodeficiency virus-infected human cytotoxic T lymphocytes during early and late TCR responses. *J Exp Med* 1994;179:111-121.

134. Kramas AJ, et al. T cell receptor usage influences specificity of human immunodeficiency virus type-1 specific cytotoxic T lymphocyte killing: analysis of dual-species recognition reveals a dominant response directed against a conserved *in vivo* variant. *J Exp Med* 1996;183:1669-1679.
135. Wilson CC, et al. Overlapping epitopes on human immunodeficiency virus type-1 gp120 prevented by HLA A, B, and C molecules: effects of viral variation on cytotoxic T lymphocyte recognition. *J Virol* 1997;71:1256-1264.
136. Wolinsky SM, et al. Adaptive evolution of human immunodeficiency virus type-1 during the natural course of infection. *Science* 1996;272:537-547.
137. Haas G, et al. Dynamics of viral variants in HIV-1: Not and specific cytotoxic T lymphocytes *in vivo*. *J Immunol* 1996;157:4217-4221.
138. Nowak MA, et al. Antigenic oscillations and shifting immunodominance in HIV-1 infections. *Nature* 1995;375:606-611.
139. Mackewicz CE, Orriya HW, Levy JA. CD8+ cell anti-HIV activity correlates with the clinical state of the infected individual. *J Clin Invest* 1991;87:1462-1466.
140. Scala E, et al. C-C chemokines, IL-16, and soluble antiviral factor activity are increased in cloned T cells from subjects with long-term nonprogressive HIV infection. *J Immunol* 1997;158:4485-4492.
141. McKenzie S, Dall'olio G, North M, Frame P, Meaux R. Serum chemokine levels in patients with non-progressing HIV infection. *AIDS* 1996;10:29-33.
142. Barusi S, D'Andrea M, Simonelli C, Tirelli G, DePaoli P. Serum levels of RANTES and MIP-1 $\alpha$  in HIV-positive long-term survivors and progressor patients. *AIDS* 1996;10:1431-1432.
143. Clerici M, et al. Chemokine production in HIV-seropositive long-term asymptomatic individuals. *AIDS* 1996;10:1432-1433.
144. Chen Y, Gupta P. CD8+ T-cell-mediated suppression of HIV-1 infection may be due to chemokines RANTES, MIP-1 $\alpha$ , and MIP-1 $\beta$ . *AIDS* 1996;10:1434-1437.
145. Blazewicz V, Hwang M, Kasper A, Luzzati T, Kohn J, RANTES, MIP and other chemokines in HIV infection. *AIDS* 1996;10:1435-1436.
146. Furtak E, et al. Antigen-enhanced chemokine-mediated HIV-1 Sialinfection by CD8+ T cells from Exposed Uninfected Individuals Expressing the HIV-1 gp120. *J Exp Med* 1997;186:455-464.
147. Sallat T, Bacon K, Fox N. Chemokine-dependent attraction of monocytes and T lymphocytes to the memory phenotype by chemokine RANTES. *Nature* 1990;347:669-671.
148. Amara A, et al. HIV co-receptor: downregulation as antiviral principle. SDF-1 dependent internalization of the chemokine receptor CXCR4 contributes to inhibition of HIV replication. *J Exp Med* 1997;186:149-156.
149. Baker M, Werner A, Bandaru V, Vignatelli K, Kurth R. HIV suppression by interleukin-1. *Nature* 1996;378:563.
150. Zwart G, et al. Antibody responses to HIV envelope and gag epitopes in HIV-1 seroconverters with rapid versus slow disease progression. *Virology* 1994;201:257-263.
151. Wong MT, et al. Longitudinal analysis of the humoral immune response to human immunodeficiency virus type-1 (HIV-1) gp160 epitopes in rapidly progressing and nonprogressing HIV-1-infected patients. *J Infect Dis* 1993;168:1527-1537.
152. Hegerl-Est E, et al. Predictors for non- and slow progression in human immunodeficiency virus (HIV) type-1 infection: low viral RNA copy numbers in serum and maintenance of high HIV-1-specific but not V3-specific antibody levels. *J Infect Dis* 1995;171:47-54.
153. Montefiori DC, et al. Neutralizing and infection-enhancing antibody responses in human immunodeficiency virus type-1 long-term nonprogressors. *Infect Dis* 1996;173:126-137.
154. Se Y, Tsang PH, Ruckwirth JC, Hwang J, Beker JG. Neutralizing antibody levels as prognostic indicators of long-term non-acquired immune deficiency syndrome (AIDS)-related disorders and survival. *J Clin Immunol* 1998;18:444-451.
155. Liang M, et al. Neutralizing antibodies against sequential androgynous human immunodeficiency virus type-1 isolates after seroconversion. *Infect Dis* 1994;170:1141-1147.
156. Ingram A, et al. Neutralizing antibody responses at various stages of infection with human immunodeficiency virus type-1. *Infect Dis* 1997 (in press).
157. McGuire M, Nielsen G, Hansen J, Pedersen C, Mathiesen E, Nielsen H. Anticognous HIV-1 neutralizing antibodies: emergence of neutralizing monoclonal virus and subsequent development of escape virus neutralizing antibodies. *J Acquir Immune Defic Syndr* 1991;4:303-307.
158. Buchbinder JC, Turner RM, Timpone J, et al. Lymphoid lesions associated with the acquired immunodeficiency syndrome. *Am J Surg Pathol* 1983;7:543-554.
159. Fernandez R, Mouradian J, Nienhuis G, Davies J. The prognostic value of histopathology in persistent generalized lymphadenopathy in homosexual men. *N Engl J Med* 1983;309:125-126.
160. Janusz G, et al. An immunohistology lab approach to persistent lymphadenopathy and its relevance to AIDS. *Clin Exp Immunol* 1985;59:257-260.
161. Scherfeld P, Clay K, Marcellet B, Hertz G, Gao R, Harper M. HTLV-II expression in infected lymph nodes and relevance to pathogenesis of lymphadenopathy. *Am J Pathol* 1986;125:436-442.
162. Rice R, Tenner BA, Kuhl J, Belli A, Kern P, Dietrich M. Spectrum of morphologic changes of lymph nodes from patients with AIDS or AIDS-related complexes. *Prog Allergy* 1996;37:81-86.
163. Palever G, Gerszten R, Vardanian Nages N, LNW. HTLV-III lymphadenitis: histological and immunohistological classification. *Scand J Immunol* 1987;28:41-49.
164. Turner R, Levine A, Gao R, Harper M, Kern P. Progressive histopathologic abnormalities in the persistent generalized lymphadenopathy syndrome. *Am J Surg Pathol* 1987;11:100-107.